

paper ID-47201607

by

Submission date: 09-Jun-2021 08:25AM (UTC-0400)

Submission ID: 1603426986

File name: paper_ID-47201607.pdf (60.14K)

Word count: 2729

Character count: 14584

Particulate Matter (PM_{2,5}) Increases MDA Levels Serum of Workers at Surabaya Bus Station

Kahar¹, Soedjajadi², Lilis Sulistyorini³

Master Program Study, Environmental Health Department, Public Health Faculty

Airlangga University, Surabaya, Indonesia¹

Environmental Health Department, Public Health Faculty, Airlangga University, Surabaya, Indonesia^{2,3}

Email: kahar.yaya22@gmail.com¹

Abstract- Particulate matter (PM_{2,5}) in the air that are free radicals and enter through the respiratory station workers. Inability of antioxidants neutralize free radicals will increase MDA levels serum and cause oxidative stress. The aim of this study was to look at the differences in MDA levels serum and PM_{2,5} influence on MDA levels serum on several main station attendant Surabaya. The design of this study was an analytic observational with cross-sectional study. The population in this study was bus station workers and administration workers, large of samples were 12 people in each group. Measurement of PM_{2,5} with EPAM 5000 and MDA levels in serum using TBA test method. The results of t test analysis 2 free samples to show that there are differences in serum MDA levels were highly significant ($p < 0.05$) in the bus station workers and administration workers. Multiple linear regression showed that there are significant PM_{2,5} exposure and body mass index ($p < 0.05$) on serum levels of MDA. The conclusion of this study is the PM_{2,5} exposure can increased serum levels of MDA on several main bus station workers. Suggestions for health officials to control the station regularly and use a mask while on duty in the area of Surabaya station.

Keyword: Bus Station, PM_{2,5}, Free Radical, Malondialdehyde, Oxidative Stress

1. INTRODUCTION

Transportation in Indonesia showed a significant rise in the number. Air pollution is composed of a pollutant that can cause health problems such communities, particulate matter, carbon monoxide (CO), ozone (O₃), nitrogen dioxide (NO₂), sulfur dioxide (SO₂). Surabaya station is one of the public places with air pollution from the exhaust emissions.

PM_{2,5} have an aerodynamic diameter of less than 2.5 μ m, is part of the more dangerous because it has the ability to penetrasi deeper into the lungs and reach the alveolar.^[1]

Constituent particles consisting of organic and non organic compounds, constituents of these can trigger the formation of ROS. Oxidant particles through the mechanism may increase seven times the activity of NF- κ B in the human respiratory tract. NF- κ B is a transcription factor that can induce genes in various proinflammatory cytokines and enzymes that result from mediators of inflammation and the immune system.^{[2][3]}

a oxygen having one unpaired electron to become reactive. high concentrations of ROS in the body will react with non-radical compounds and will spur the emergence of such a chain reaction of peroxide fat. These reactive oxygen species can also cause damage to the body of molecules, including proteins, carbohydrates and DNA. Due to the high possibility of damage caused by the biological systems that are radical oxygen compounds are considered to

have contributed to aging factors and the incidence of other chronic diseases.^{[4][5]}

Oxidative stress occurs when the ROS is not sufficiently neutralized by antioxidants such as superoxide dismutase (SOD), Catalase (Cat), Glutathion Peroxide (GPx). Malondialdehyde (MDA) is an indicator of lipid peroxides. The ratio of MDA can be used as an indicator of oxidative stress were measured in serum using the TBA test. If the value of MDA increased, then this indicates the high oxidative stress conditions.^[6]

Serum MDA activity is influenced by characteristics such officer, age, smoking habits and body mass index (BMI). The aim of this study was to determine the effect PM_{2,5} on MDA levels serum on workers at Surabaya bus station.

2. METHOD

The design of this study was an analytic observational research with cross-sectional study. The population in this study consisted of two groups, bus station workers and administration workers with size sample 12 people in each group with formula.^[17]

$$n = \frac{2 \cdot \sigma^2 \cdot (Z_{\alpha} + Z_{\beta})^2}{(\mu_1 - \mu_2)^2}$$

Spesification:

Z_{α} = adjusted standart deviation α (1,96)

Z_{β} = adjusted standart deviation β (0,84)

σ = SD response to the comparison group (0,81)

μ_1 = the mean level of sample group (4,28)

μ_2 = the mean of comparison group (3,34)

PM_{2,5} were measured by Haz-Dust EPAM 5000 particulate air monitoring. MDA levels used TBA test method with spectrophotometric.

Data analysis using Mann-Whitney statistical program tests to show the differences in levels of serum and multiple linear regression to determine the effect PM_{2,5} to MDA level serum

3. RESULT

Respondents in this study were bus station workers and administration workers with the criteria was male, has worked at least one year, did not suffer from diabetes. Characteristics of workers can be seen in Table 1.

Table 1. Characteristic Bus Satation and Administration Workers

Characteristic	Bus Station Workers		Administration Workers	
	n	%	n	%
Age (Years)				
21-30	0	0	7	29,2
31-40	1	4,2	5	20,8
>40	11	45,8	0	0
mean±SD	45,83±6,04		30,25±6,28	
Smoking Habits				
No smoking (0)	5	20,8	2	8,3
Rarely (1-200)	2	8,3	9	37,5
Medium (201-600)	5	20,8	1	4,2
mean±SD	164,8±171,3		50,3±71,1	
Mass Body Index				
Less (<18,4)	0	0	2	8,3
Normaly (18,5-25,0)	3	12,5	7	29,2
Obesity (>25,1)	9	37,5	3	12,5
mean±SD	28±4,33		22,43±3,58	

Table 1 shows that workers in the station with a mean age of 45.83 years, classified as light smokers (Index Briskman/IB) with a mean of 164.8 cigarettes/year and with a mean body mass index of 28kg/cm², including obesity. whereas the clerical workers 30.25 years, including light smokers of 50.3 cigarettes/year and a body mass index of 22.4 kg/cm² was normal.

Table 2. PM_{2,5} levels in the bus station and an administration

Level Measurement	Bus Station	Administration room
PM _{2,5}	0,82 mg/m ³	0,11 mg/m ³
Temperature	33°C	29,3°C
Humadity	57,8%	67%

The results of PM_{2,5} measured that the concentrations of PM_{2,5} in Station showed a mean value of 0.829 mg/m³. While the administration room obtained a mean value of 0.09 mg/m³. PM_{2,5} levels in Station above the threshold value. Based on the Decree of the Minister of Health of the Republic of Indonesia No. 1405 / Menkes / SK / XI / 2002 regarding Requirements Environmental Health Office Work and Industry at 0.15 mg/m³^[17].

Levels of malondialdehyde (MDA) is a marker of oxidative stress by checking TBA test using spectrophotometric method. MDA level examination results are shown in Table 3:

Table 3. Results of Measurements of Serum Levels of MDA in Bus Station Workers and Administration workers

Malondialdehyde (MDA) Levels (nmol/ml)		
Workers	mean±SD	ρ
Bus station	9,44±2,94	0,01**
Administration	5,60±0,60	

** ρ <0,05= significant

Table 3 shows that the MDA levels serum in bus station workers higher than administration workers. It was caused by bus station workers exposed PM_{2,5} which one pollutant that is free radicals. There are very significant differences bus station workers and administration workers (ρ <0.05).

Table 4. Analysis of The Effect of PM_{2,5} Exposure to Increases MDA Levels Serum in Bus Station Workers and Administration Workers.

Variabel	MDA Level	
	β	ρ
Particulate Matter (PM _{2,5})	1,143	0,046*
Age	0,313	0,164
Smoking Habits	0,142	0,575
Mass Body Index	0,543	0,028*

* ρ <0,05= significant

Based on the results of multiple linear regression analysis showed that PM_{2,5} affect MDA (β = 1.143; ρ = 0.046). PM_{2,5} exposed higher in the air can increases MDA levels serum. Cigarette smoke contains components of gases and particles that potential to cause free radical-induced oxidative stress

in the body, lipid peroxidation (MDA) and lower levels of endogenous antioxidants.^[8]

Body Mass Index (BMI) ($\beta = 0.543$; $\rho = 0.028$). Workers in the station majority obesity. This study is in line with Selvakumar (2012)^[9], shows that there is the influence of obesity with increased activity of MDA ($\beta = 0.483$; $\rho = 0.003$). Obesity not only associated with being overweight but always associated with type 2 diabetes metabolic disorders due to excess accumulation calories into fat and saturated fat release to various organs with high concentrations will cause chronic oxidative stress^[10]^[17]

4. DISCUSSION

PM_{2.5} in the air can increase serum levels of MDA. Particles enter the body primarily through the respiratory system due to the direct adverse effect mainly occurs in the respiratory system. Particles larger than 5 microns will gather and stop especially on the nose and throat. 0.5-5 micron-size particles can gather in the lungs to reach bronchioli and only a small portion can only be issued by the cilia in about 2 hours. While the particle size of 0.5 microns will reach and gather in the alveoli. Activities Cleaning particles are very small yag of alveoli running very slow and imperfect when compared with activity in the larger channel. Even some particles exist which still remained in the alveoli and absorbed in the blood^[11].

Lipid peroxidation can be described generally as a process in which oxidants such as free radicals attack the lipids containing double carbon, especially polyunsaturated fatty acids, especially (PUFAS / Polyunsaturated Fatty Acids). One of the consequences of uncontrolled oxidative stress (an imbalance between prooxidant and antioxidant levels in support prooxidants) are cells, tissues, and organs caused by oxidative damage. A high level of free radicals or reactive oxygen species (ROS) can cause direct damage to lipids. The main source of endogenous ROS production is mitochondria, plasma membrane, endoplasmic reticulum, peroxisomes and through various mechanisms, including enzymatic reactions or autooxidation of some compounds, such as catecholamines and hydroquinone. Different exogenous stimuli, such as ionizing radiation, ultraviolet rays, tobacco smoke, pathogen infection, environmental toxins, and exposure to herbicides / insecticides, is the source of the in vivo production of ROS^[12].

Components of chemicals that exposure of air pollutants that cause oxidative stress and inflammation can be in the form of gases and particles. Contained in the oxidative potential of particles depend on particle composition, size, especially the ultrafine particle concentrations and other transition metals and organic

chemicals such as semi-volatile and volatile (Delfino, 2011). reactive chemicals that are important can include organic components which themselves PM_{2.5} as quinones oxidize or PAH species that undergo biotransformation by cytochrome P-450 1A1 to quinones, which can induce oxidative stress (Bonvallot, 2001).

Body Mass Index has been found to increase levels of serum MDA. ROS pathways in mitochondria increased in obesity. Within 24 hours it takes at least as much as 352.81 liters of oxygen. The purpose is filled with breathing less than 23 thousand times. The consequences of which are from the metabolic processes in biochemical systems (biological oxidation) in the body is able to produce free radicals as much as 2.5% of the total oxygen requirement 3,4kg / 24 hours. Though the biological oxidation can take place without oxygen, but all mammals requires absolute oxygen supply through breathing (respiration). Respiration is the process of formation of adenosine triphosphate (ATP) as an energy, which is obtained from a reaction between hydrogen and oxygen that then form water. Reaction energy formation known as oxidative phosphorylation takes place in the mitochondria. The process of respiration takes place in the mitochondria maatriks through a reaction which is then referred to as the respiratory chain. Thus, the formation of ROS within mitochondria is triggered by chronic electron leakage from normal respiratory chain. So, on the chest wall will decrease obesity and respiratory work will increase (Widayati, 2012; Limanan, 2013).

MDA is the end product produced by the decomposition of arachidonic acid and polyunsaturated fatty acids (PUFAS) greater peroxidation in the cell. The increase in free radicals causing overproduction of MDA. Malondialdehyde levels commonly known as a marker of oxidative stress and antioxidant status (Gawel, 2003).

5. CONCLUSION

PM_{2.5} concentration in several main bus station is higher than in the administration is. PM_{2.5} high and obesity was effects MDA levels serum in the bus station workers and administrative workers. MDA levels serum were high indicates oxidative stress in bus station workers.

6. SUGGESTION

Air quality monitoring on a regular basis to controlling air quality around the bus station. Workers in the station should be health checked regularly as early prevention and consume fruits and vegetables that contain lots of antioxidants to support the enzymatic antioxidants in the body.

Acknowledgement

Thanks to several main bus station workers at Surabaya, which was willing to become respondents in this study.

REFERENCES

- [1] WHO. 2005. *WHO Air Quality Guidelines for Particulate Matter, Ozone, Nitrogen Dioxide and Sulfur Dioxide*. WHO Regional Office for Europe. Germany: Druckpartner Moser
- [2] Li N, Xia T and Andre EN. 2008. The Role of Oxidative Stress in Ambient Particulate Matter-Induced Lung Disease and Its Implication in The Toxicity of Engineered Nanoparticles. *Free Radical Biology and Medicine*. 44 (9): 1689-1699
- [3] Yang, W., and Omaye, S. T. 2009. Air pollutants, oxidative stress and human health. *Mutation Research/Genetic Toxicology and Environmental Mutagenesis*, 674 (1): 45-54.
- [4] Papas A.M., 1996. Determinants of antioxidant status in humans. *Lipids*, 31 (1): S77-S82.
- [5] Lodovici M and Elisabetta B. 2011. Review Article: Oxidative Stress and Air Pollutant Exposure. *Journal of Toxicology*: 1-9
- [6] Jyothi P., Riyaz N., Nandakumar G., Ninitha MP. 2008. A Study of Oxidative Stress in Paucibacillary and Multibacillary Leprosy. *Indian Journal of Dermatology, Venereology and Leprology*, 75 (1): 80Hulley, 2006
- [7] Hulley SB, Cummings SR, Browner WS, Grady D, Heerst N, Newman TB. 2006. *Designing Clinical Research: An Epidemiological Approach*. 3rd Ed. Baltimore: Lippincott Williams and Wilkins
- [8] Agnihotri R, Pandurang P, Kamath SU, Goyal R, Ballal S, Shanbhogue AY, Kamath U, Bhat GS, and Bhat KM. 2009. Association of Cigarette Smoking with Superoxide Dismutase Enzyme Level in Subjects with Chronic Periodontitis. *Journal Periodontol*. 80 (4): 657-662
- [9] Selvakumar, C., and Uma, M. 2012. Oxidant-antioxidant disturbance in men classified as obese according to the preliminary WHO guidelines for Asians. *Journal of Stress Physiology and Biochemistry*, 8 (1): 172-181Abdali D., Samson SE and Grovek A.K. 2015. How Effective are Anioxidanta Supelements in Obesity and Diabetes?. *Medical Principle and Practice*. 24 (3): 201-215
- [10] Valdecantos, M. P., Pérez-Matute, P., and Martinez, J. A. 2008. [Obesity and oxidative stress: role of antioxidant supplementation]. *Revista de investigacion clinica; organo del Hospital de Enfermedades de la Nutricion*, 61 (2): 127-139.
- [11] Florence T., Beatriz GF and Lester K. 2003. Serial Review : Role of Reactive Oxygen and Nitrogen Species (ROS/RNS) in Lung Injury and Disease. *Free Radical Biology and Medicine*. 35 (4): 327-340
- [12] Ayala A., Muñoz M. F., and Argüelles, S. 2014. Lipid peroxidation: production, metabolism, and signaling mechanisms of malondialdehyde and 4-hydroxy-2-nonenal. *Oxidative medicine and cellular longevity*, (360438), 1-32
- [13] Bonvalot V., Baeza-Squiban A., Baulig A., Brulant S., Boland S., Muzeau F., ... and Marano, F. 2001. Organic compounds from diesel exhaust particles elicit a proinflammatory response in human airway epithelial cells and induce cytochrome p450 1A1 expression. *American journal of respiratory cell and molecular biology*, 25(4): 515-521.
- [14] Widayati, E. 2012. Oxidasi Biologi, Radikal Bebas, dan Antioxidant. *Majalah Ilmiah Sultan Agung*, 50 (128): 26-32
- [15] Limanan, D. 2013. Hantaran Sinyal Leptin dan Obesitas: Hubungannya dengan Penyakit Kardiovaskuler. *eJurnal Kedokteran Indonesia*, 1 (2): 144-155.
- [16] Gawel S., Wardas M., Niedworok E., and Wardas, P. 2003. Malondialdehyde (MDA) as a lipid peroxidation marker. *Wiadomosci lekarskie (Warsaw, Poland: 1960)*, 57 (9-10): 453-455.
- [17] Abdali D., Samson SE and Grovek A.K. 2015. How Effective are Anioxidanta Supelements in Obesity and Diabetes?. *Medical Principle and Practice*. 24 (3): 201-215.
- [18] Delfino R.J., Staimer N., and Vaziri N.D. 2011. Air pollution and circulating biomarkers of oxidative stress. *Air Quality, Atmosphere and Health*, 4(1): 37-52.
- [19] Minister of Health of the Republic of Indonesia No. 1405 / Menkes / SK / XI / 2002 regarding Requirements Environmental Health Office Work and Industry

ORIGINALITY REPORT

12%

SIMILARITY INDEX

10%

INTERNET SOURCES

7%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1

Submitted to American International School -
Chennai

Student Paper

1%

2

G Istifanus, O Grace, W Luka, A James.
"Comparative Study on the Effects of Locally
and Industrially Processed Olive Oil on Some
Selected Organs of Albino Wister Rats",
Journal of Applied Life Sciences International,
2017

Publication

1%

3

Submitted to New University of Astana

Student Paper

1%

4

journals.plos.org

Internet Source

1%

5

Meigen Liu, Kimitaka Hase, Yoshihisa
Masakado, Tetsuya Tsuji, Yohei Otaka, Naoichi
Chino. "Rehabilitation research from a non-
English speaking country published in peer-
reviewed English journals – the Japanese
experience", Disability and Rehabilitation,
2009

1%

6	Rocío de la Iglesia. "Healthy properties of proanthocyanidins", BioFactors, 03/15/2010 Publication	1 %
7	itfx.pitstop-ng.com Internet Source	1 %
8	Alex Woodell, Bärbel Rohrer. "Chapter 38 A Mechanistic Review of Cigarette Smoke and Age-Related Macular Degeneration", Springer Science and Business Media LLC, 2014 Publication	1 %
9	Submitted to UC, Boulder Student Paper	1 %
10	webgate.ec.europa.eu Internet Source	1 %
11	Kasım Takım, Mesut Işık. "Phytochemical Analysis of Paliurus spina-christi Fruit and Its Effects on Oxidative Stress and Antioxidant Enzymes in Streptozotocin-Induced Diabetic Rats", Applied Biochemistry and Biotechnology, 2020 Publication	1 %
12	theses.uin-malang.ac.id Internet Source	1 %
13	ejournals.epublishing.ekt.gr Internet Source	1 %

14	icvl.ru Internet Source	1 %
15	text-id.123dok.com Internet Source	<1 %
16	Ernawati, Erry Gumilar, Kuntoro, Joewono Soeroso, Gus Dekker. "Expectant management of preterm preeclampsia in Indonesia and the role of steroids", The Journal of Maternal-Fetal & Neonatal Medicine, 2015 Publication	<1 %
17	www.i-scholar.in Internet Source	<1 %
18	www.lexisnexis.co.za Internet Source	<1 %
19	Submitted to Universitas Diponegoro Student Paper	<1 %
20	pdfs.semanticscholar.org Internet Source	<1 %
21	epdf.tips Internet Source	<1 %
22	ojs.ukw.edu.pl Internet Source	<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography Off