

# CHEMICAL COMPOSITION AND SOURCES ATTRIBUTION OF RAINWATER IN BANDUNG AREA, INDONESIA

*by* Nia Yuniarti Hasan, Driejana, Aminudin Sulaeman, Herto Dwi  
Ariesyady

---

**Submission date:** 09-Jun-2023 12:12PM (UTC+0700)

**Submission ID:** 2112272618

**File name:** GEOMATE\_NIA\_YUNIARTI\_HASAN.pdf (396.52K)

**Word count:** 4454

**Character count:** 23051

## CHEMICAL COMPOSITION AND SOURCES ATTRIBUTION OF RAINWATER IN BANDUNG AREA, INDONESIA

Nia Yuniarti Hasan<sup>1</sup>, \*Driejana<sup>2</sup>, Aminudin Sulaeman<sup>3</sup>, and Herto Dwi Ariesyady<sup>4</sup>

<sup>1</sup>Environmental Engineering Doctoral Program, Faculty of Civil and Environmental Engineering, Institut Teknologi Bandung, Indonesia, <sup>2</sup>Faculty of Civil and Environmental Engineering, Institut Teknologi Bandung, Indonesia, and <sup>3</sup>Faculty of Mathematics and Natural Science, Institut Teknologi Bandung, Indonesia,

\*Corresponding Author, Received: 20 May 2019, Revised: 13 Aug. 2019, Accepted: 28 Aug. 2019

**ABSTRACT:** Data of international and national monitoring stations have showed evidences of rainwater acidification in Indonesia. This research aimed to investigate the small-scale variations and the influence of urban activities to rainwater chemical composition in Bandung, Indonesia, as well as to determine the contributing sources to its chemical variation. Rainwater bulk samples had been collected weekly at (four) locations representing differences in altitude and local land use, from February to September 2016. Samples were analyzed for pH, conductivity, and major ions ( $\text{SO}_4^{2-}$ ,  $\text{NO}_3^-$ ,  $\text{Cl}^-$ ,  $\text{Na}^+$ ,  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$ ,  $\text{K}^+$ , and  $\text{NH}_4^+$ ). The results showed that the pH of rainwater were in the range of acidic to normal (average  $5.42 \pm 0.72$ ), where ammonium acted as the major neutralizing factor (NF). Small-scale spatial variability of  $\text{nss-SO}_4^{2-}$ ,  $\text{NO}_3^{2-}$ ,  $\text{nss-Cl}^-$ ,  $\text{nss-Ca}^{2+}$ ,  $\text{nss-Mg}^{2+}$ , and  $\text{NH}_4^+$  within 4 (four) locations were observed. Sulphate and  $\text{NO}_3^-$  as well as  $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$  were significantly correlated, with  $r = 0.870$  and  $r = 0.929$ , respectively. Principal Component Analysis (PCA) indicated three main sources of rainwater chemistry. Firstly, a mix of natural sources (e.g., sea-salt aerosols, crustal elements and waste decomposition) which consisted of sea-salt  $\text{SO}_4^{2-}$ ,  $\text{ss-Cl}^-$ ,  $\text{Na}^+$ ,  $\text{ss-Ca}^{2+}$ ,  $\text{nss-Ca}^{2+}$ ,  $\text{ss-K}^+$ ,  $\text{nss-K}^+$  and  $\text{NH}_4^+$ . Secondly, fossil fuel combustion/vehicular emissions, which consisted of  $\text{nss-SO}_4^{2-}$ ,  $\text{NO}_3^-$ ,  $\text{H}^+$ , and  $\text{nss-Mg}^{2+}$ ; and thirdly,  $\text{nss-Cl}^-$  presumably from other acid reaction. These three components account for 80.25% of the total variance. The results suggested that local and long-range transported emissions of natural and anthropogenic sources contributed to the rainwater chemistry in the Bandung area.

*Keywords:* Acid Rain, Wet Deposition, PCA, Rainwater Chemistry

### 1. INTRODUCTION

Chemical compositions of rainwater have been studied in many urban areas worldwide, such as to evaluate the impact of air pollution. Source analysis and contribution of anthropogenic and natural sources to rainwater contamination are important in the context of air quality management [1,2,3,4,5,6,7,8,9,10,11,12,13,14,15,16,17,18].

The pH of rainwater could be used as an indicator of atmospheric pollution because under normal conditions, based on the value in the equilibrium of pure water and atmospheric  $\text{CO}_2$ , pH of rainwater is 5.6 [19]. Acidity variability occurs due to atmospheric chemistry processes involving air pollutants absorbed from various sources and being washed off by rain droplets. For examples, the alkaline pH in rainwater in the Pune Region of India mainly caused by the abundance of  $\text{Ca}^{2+}$  originated from vehicle driven roadside dust [2]. Acidic rainwater in Shihwa Basin Korea was influenced by the dominant ions of  $\text{SO}_4^{2-}$  and  $\text{NO}_3^-$  which were neutralized by  $\text{NH}_4^+$  and  $\text{Ca}^{2+}$  [17].

Elevation differences and local anthropogenic

activities might account for the variation in the rainwater chemical composition [13], causing, e.g. monthly variations of rainwater chemistry in the Yulong Mountain Region in Southwestern China.

Bandung area is located in a mountainous region of West Java Indonesia where the lowest level is at an elevation of 768 meters above sea level (asl). Bandung is the third most populous city in Indonesia after Jakarta and Surabaya. It is the capital of the West Java Province, located about 140 km from Jakarta as the national capital. Its urban activities potentially affect its air quality, and hence rainwater, to some extent. Bandung has a tropical, wet, and dry climate with average temperature ranges from 18.1°C to 31.9°C during 2015 and an average rainfall of 184.75 mm with the ranges of 0.3 mm/month (in July during the dry season) to 322.4 mm/month (in November in the wet season).

Due to its characteristics, rainwater quality in the Bandung area might be influenced by air pollution. This paper explores rainwater characteristics in Bandung and identifies the likely sources that determine its chemical composition. Samples were collected during the rainy events from February to September 2016.

29

## 2. METHODOLOGY

### 2.1 Sampling Area

Rainwater samples were collected at 4 (four) locations, representing different elevation and land use within the Bandung area, as shown in Figure 1 as follow :

- 1) Lembang (06°49'35,01" S 107°37'04,54" E) with an altitude of approximately 1.272 m asl lies on the hillside of the mountain area. The site is located approximately 12 km away from the center of Bandung City, with the land use predominantly being agricultural, livestock, and tourism activity.
- 2) Coblong (06°55'11,07" S 107°36'04,54" E) with an altitude of approximately 803 m asl. It is an urban area with high traffic activity.
- 3) Sumur Bandung (06°55'11,07" S 107°36'39,34" E) with an altitude of approximately 742 m asl. It is the center of major urban activity such as offices complex, trading, and traffic area.
- 4) Buah Batu (06°53'26,69" S 107°36'43,73" U) with an altitude of approximately 642 m asl is a sampling site in urban settlement area with high population density and their associated activities.



Fig. 1. Sampling Site

26

### 2.2 Samples Collection and Analysis

Rainwater samples were taken on a weekly basis from February to September 2016. The rainwater samples were collected using a polyethylene bottle (1.000 mL). Rainwater passed through a polyethylene funnel (12 mm in diameter) equipped with a 1 mm filter plug that was put inside a stainless steel cylinder. The sampler bottles, funnels and filters were cleaned and rinsed with deionized water before each use. Rainwater samples were then transferred to 100 mL PE bottle for ion analysis.

The pH values and conductivity for the

rainwater samples were analyzed with a pH meter (Sartorius Professional Meter PP-20), and a conductivity meter (Agilent Technologies 3200 C). Major ions in rainwater of sulphate ( $\text{SO}_4^{2-}$ ), nitrate ( $\text{NO}_3^-$ ), chloride ( $\text{Cl}^-$ ), calcium ( $\text{Ca}^{2+}$ ), magnesium ( $\text{Mg}^{2+}$ ), sodium ( $\text{Na}^+$ ), potassium ( $\text{K}^+$ ), and ammonium ( $\text{NH}_4^+$ ) were determined using Ion Chromatography (DIONEX ICS 500 DP).

Statistical characteristics of rainfall, pH, conductivity, and Precipitation-Weighted Mean (PWM) of ion concentration were analyzed. The spatial variation between the sampling sites was analyzed using One-way ANOVA. Source attribution was analyzed using correlation and Principal Component Analysis (PCA).

34

### 2.1 Quality Control of Samples Analysis

The quality control of the samples was checked by Ion Balance (IB) defined as Ion Percent Difference (IPD), as  $\text{IPD} = [(C - A) / (C + A)] \times 100$ , where, C and A represents anion and cation equivalents ( $\mu\text{eq L}^{-1}$ ), respectively [20]. Precipitation weighted mean (PWM) was carried out by calculating the sum of the product of ion concentration and the precipitation amount, obtained from rainwater sample, for each data pair, and dividing the result by the sum of precipitation amount for a certain period [21].

30

## 3. RESULT AND DISCUSSION

### 3.1 Rainfall Amount, pH and Conductivity of Rainwater

Rainfall amount at 4 (four) sampling sites was largely varied, ranging between 40.69 – 347.66 mm/month. High rainfall events during 2016 were suggested to be impacted by the La Nina event (Fig.2), with the highest events recorded in March, April, and May.




Fig.2 Rainfall Distribution at Four Sampling Sites

36

The pH of the rainwater ranged from 4.74 to 6.20 (average 5.45), lower than the pH of typical natural rainwater (5.6), indicating the influence of acidic substances in the rainwater [19].

Rainwater from Buah Batu showed a tendency of more acidic compared to other sites, particularly the Lembang site which is located in the rural area at the highest altitude. This condition is contrary to the acidic orographic enhancement found in Northern England [23], suggesting local urban emission sources might have a more significant influence in the chemistry of rainfall.

Nearly 60% of acid precipitation pH was between 4.5 to less than 5.6, while 42% had pH less than 5.0 (which would have significant effects on environmental acidification). Within this 42%, 11% were strongly acidic with pH < 4.5 (Fig. 3).




Fig.3 Rainwater pH Distribution

The major precursors of acidity in rainwater were SO<sub>2</sub> and NO<sub>x</sub> as were found in the sources attribution analysis (See Section 3.5). Soil particles and sea spray origin species might play an essential role in neutralizing SO<sub>2</sub> and NO<sub>x</sub>. The abundance of acid precursors is seen in the rainwater conductivity, the value being related to the total sum of anions and cations. Rainwater conductivity is mainly contributed by water-soluble ions. The rainwater electric conductivity ranged from <0.5 to 100 μS/cm (Fig. 4). There is no clear pattern as to whether conductivity was related to the rainwater amount, as the temporal variation of conductivity did not resemble the pattern of rainfall amount as seen in Fig. 2.




Fig.4 Temporal Variation of Conductivity (μS/cm)

### 3.2 Temporal Variation of Precipitation-Weighted Mean (PWM) Concentration

Precipitation-Weighted Mean (PWM) concentrations were calculated to eliminate the dilution effect of rainwater quantity during high rainfall. The temporal variations of PWM concentration of major ions that reflected the variation due to acid chemistry processes in rainwater are shown in Figure 5.

While rainfall monthly variation showed clear pattern of seasonal influence of lowest amount during June – August at all sites, SO<sub>4</sub><sup>2-</sup> and NO<sub>3</sub><sup>-</sup> PWM concentrations showed irregular patterns. Particularly for SO<sub>4</sub><sup>2-</sup> and NO<sub>3</sub><sup>-</sup>, the PWM concentrations were detected during low and high rainfall, suggesting that rainfall was not the sole factor affecting the concentration. Instead, the variation might be due to the strength of acid precursor emissions from local and regional sources. On the other hand, increasing ammonium (NH<sub>4</sub><sup>+</sup>) as well as Cl<sup>-</sup>, Na<sup>+</sup>, Ca<sup>2+</sup>, Mg<sup>2+</sup>, and K<sup>+</sup> ions showed in the PWM concentration on lower rainfall suggested the dilution effect of these natural-origin ions. In addition, the strong temporal variation of anion SO<sub>4</sub><sup>2-</sup> and cation NH<sub>4</sub><sup>+</sup> were presumably caused by the neutralization reaction in the rainout and washout deposition processes.




Fig.5 Temporal Variation of Precipitation Weight Mean (PWM) Concentration Ions

### 3.3 Neutralizing Factors

Neutralizing factor (NF) was used to evaluate the neutralization of major anions by crustal components ( $\text{Ca}^{2+}$  and  $\text{Mg}^{2+}$ ) and  $\text{NH}_4^+$ . The neutralization of acids by base cations were evaluated by calculating the neutralization factors using the formula:  $\text{NF}(x) = ((X^+) / (\text{SO}_4^{2-} + \text{NO}_3^-))$  where X is the  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{NH}_4^+$ .

The NF observed for  $\text{Ca}^{2+}$ ,  $\text{Mg}^{2+}$  and  $\text{NH}_4^+$  were 0.31, 0.09, 0.56 (Lembang); 0.26, 0.05, 0.56 (Coblong); 0.32, 0.05, 0.49 (Sumur Bandung) and 0.38, 0.06, 0.43 (Buah Batu). Hence, NF showed that  $\text{NH}_4^+$  was the dominant neutralizing ion in the Bandung area.

This result was similar to what was found in Chiang May Thailand [1], Dalian China [6], Ya'an, Southwestern China [9], and Nanpin Mangdang Mountain China [25].

### 3.4 Sea salt and Non-Sea Salt Contribution

PWM concentration of sea salt (ss) and non-sea salt (nss) contributions were described in Figure 6. Data used for analyzing ss-ions and nss-ions were the long-term monitoring by EANET (Acid Deposition Monitoring Network in East Asia). Non-sea salt  $\text{SO}_4^{2-}$  and nss- $\text{Ca}^{2+}$  were the dominant ions for all sampling sites. Non-sea salt sulfate was the highest anion PWM concentration in the Bandung area for more than ten years of EANET monitoring. The trend of  $\text{SO}_2$  emission during the 2000 – 2014 period in Indonesia increased by 50% ( $+ 0.12 \text{ ppb year}^{-1}$ ) [23]. The observation found in this study was in line with the EANET finding.




Fig.6 PWM Concentration of Sea Salt and Non-Sea Salt Contribution

### 3.5 Sources Attribution Analysis

#### 3.5.1 One-Way ANOVA

One-way analysis of variance (ANOVA) was used to analyze the spatial variability of the ions in rainwater chemistry, as shown in Table 1. Statistical analysis was applied to log-normal transformed data because the raw data of rainwater amount and ion concentrations had positive skewness distribution.

The F-test statistic was applied to test the null hypothesis; that there is no variance of ions concentration and rainfall amount between different sampling sites (0.05).

It was found that the concentration of nss- $\text{SO}_4^{2-}$ ,  $\text{NO}_3^{2-}$ , nss- $\text{Ca}^{2+}$ , nss- $\text{Mg}^{2+}$ , and  $\text{NH}_4^+$  ions were significantly different between the 4 (four) sampling sites indicating small-scale spatial

variability.

Table 1 ANOVA of Ion Concentration in Rainfall and Rainfall Amount

	F	Sig.
Rainfall (mm)	1.289	0.281
ss-SO <sub>4</sub> <sup>2-</sup>	0.252	0.860
nss-SO <sub>4</sub> <sup>2-</sup>	<b>11.353</b>	<b>0.000</b>
NO <sub>3</sub> <sup>-</sup>	<b>3.258</b>	<b>0.024</b>
ss-Cl <sup>-</sup>	0.076	0.973
nss-Cl <sup>-</sup>	1.104	0.351
H <sup>+</sup>	0.479	0.698
Na <sup>+</sup>	0.252	0.859
ss-Ca <sup>2+</sup>	0.252	0.860
nss-Ca <sup>2+</sup>	<b>7.991</b>	<b>0.000</b>
ss-Mg <sup>2+</sup>	0.088	0.967
nss-Mg <sup>2+</sup>	<b>2.931</b>	<b>0.037</b>
ss-K <sup>+</sup>	0.253	0.859
nss-K <sup>+</sup>	0.517	0.671
NH <sub>4</sub> <sup>+</sup>	<b>2.932</b>	<b>0.037</b>

3.5.2 Ion Correlation

Correlation between ions was used to determine the different sources of anions and cations in the rainwater and their potential linkages. Ions concentration in rainwater was affected by rainfall amount, possibly causing a multicollinearity problem. Partial correlation analysis of the strength and direction of a linear relationship between ions whilst controlling for rainwater amount were calculated and presented in

Table 2.

The acidic ions of non-sea salt SO<sub>4</sub><sup>2-</sup> and NO<sub>3</sub><sup>-</sup> that significantly correlated (r = 0.901) possibly indicate similar sources. Natural rainwater becomes more acidic from the sulfuric and nitric acids that are formed by acidic gases of SO<sub>2</sub> and NO<sub>3</sub> from anthropogenic activities.

Significant correlation between SO<sub>4</sub><sup>2-</sup> and NH<sub>4</sub><sup>+</sup> (r = 0.431) also NO<sub>3</sub><sup>-</sup> and NH<sub>4</sub><sup>+</sup> (r = 0.481) indicated that ammonium exists in the atmosphere as (NH<sub>4</sub>)<sub>2</sub>SO<sub>4</sub> and NH<sub>4</sub>NO<sub>3</sub> [19]. On the other hand, a significant correlation between Ca<sup>2+</sup> and Mg<sup>2+</sup> (r = 0.801) showed their occurrence from the crustal origin.

3.5.3 Principal Component Analysis (PCA)

Principal components analysis (PCA) was used to identify and associate potential sources distribution of the species in rainwater chemistry. PCA was applied to explore data relationships and source types for rainwater composition in many studies [1, 2, 5, 9, 10, 11, 13, 16, 18]. As a cut-off, factors with an eigenvalue >1 were retained. Factors are extracted to be interpreted as a common source for the components in the same factor. In this study, PCA explained 82.291% of the data variance.

Table 2. Partial Correlation Coefficients of Major Ions in Rainwater

	nssSO <sub>4</sub> <sup>2-</sup>	NO <sub>3</sub> <sup>-</sup>	ssCl <sup>-</sup>	nssCl <sup>-</sup>	H <sup>+</sup>	Na <sup>+</sup>	ssCa <sup>2+</sup>	nssCa <sup>2+</sup>	ssMg <sup>2+</sup>	nssMg <sup>2+</sup>	ssK <sup>+</sup>	nssK <sup>+</sup>	NH <sub>4</sub> <sup>+</sup>
ssSO <sub>4</sub> <sup>2-</sup>	.563	.530	1.000	-.197	.109	1.000	1.000	.619	1.000	.449	1.000	.617	.430
nssSO <sub>4</sub> <sup>2-</sup>	.000	.000	.000	.055	.291	.000	.000	.000	.000	.000	.000	.000	.000
NO <sub>3</sub> <sup>-</sup>		.901	.563	.079	.178	.563	.563	.780	.563	.611	.562	.389	.431
ssCl <sup>-</sup>		.000	.000	.446	.084	.000	.000	.000	.000	.000	.000	.000	.000
nssCl <sup>-</sup>			.530	.009	.226	.530	.530	.627	.530	.438	.530	.366	.481
H <sup>+</sup>			.000	.929	.027	.000	.000	.000	.000	.000	.000	.000	.000
Na <sup>+</sup>				-.198	.109	1.000	1.000	.619	1.000	.449	1.000	.616	.430
ssCa <sup>2+</sup>				.053	.290	.000	.000	.000	.000	.000	.000	.000	.000
nssCa <sup>2+</sup>					-.009	-.197	-.196	.048	-.197	.124	-.197	.107	-.095
ssMg <sup>2+</sup>					.930	.055	.055	.646	.055	.227	.054	.300	.358
nssMg <sup>2+</sup>						.109	.110	.062	.109	.041	.109	.075	-.154
ssK <sup>+</sup>						.291	.288	.549	.290	.691	.292	.470	.134
nssK <sup>+</sup>							1.000	.619	1.000	.449	1.000	.617	.430
NH <sub>4</sub> <sup>+</sup>							.000	.000	.000	.000	.000	.000	.000
								.619	1.000	.450	1.000	.617	.430
								.000	.000	.000	.000	.000	.000
									.619	.801	.618	.440	.282
									.000	.000	.000	.000	.005
										.449	1.000	.617	.430
										.000	.000	.000	.000
											.449	.379	.292
											.000	.000	.004
												.616	.430
												.000	.000
													.204
													.046

Table 3. Component Matrix of the Sources Attribution Analysis using Principal Component Analysis (PCA).

	Component		
	1	2	3
ss-SO <sub>4</sub> <sup>2-</sup>	<b>0.955</b>	0.272	0.015
nss-SO <sub>4</sub> <sup>2-</sup>	0.304	<b>0.900</b>	0.080
NO <sub>3</sub> <sup>-</sup>	0.355	<b>0.853</b>	-0.008
ss-Cl <sup>-</sup>	<b>0.955</b>	0.272	0.015
nss-Cl <sup>-</sup>	-0.179	0.151	<b>0.822</b>
H <sup>+</sup>	-0.365	<b>0.402</b>	-0.518
Na <sup>+</sup>	<b>0.955</b>	0.272	0.015
ss-Ca <sup>2+</sup>	<b>0.955</b>	0.272	0.015
nss-Ca <sup>2+</sup>	<b>0.639</b>	0.548	0.341
ss-Mg <sup>2+</sup>	<b>0.955</b>	0.272	0.015
nss-Mg <sup>2+</sup>	<b>0.479</b>	<b>0.522</b>	0.427
ss-K <sup>+</sup>	<b>0.955</b>	0.271	0.14
nss-K <sup>+</sup>	<b>0.760</b>	0.168	0.326
NH <sub>4</sub> <sup>+</sup>	<b>0.430</b>	0.376	-0.008
Total Variance (%)	<b>62.750</b>	<b>10.780</b>	<b>8.761</b>

Factor 1 explained approximately 60.75 % of the total variance, with strong positive loadings for sea salt SO<sub>4</sub><sup>2-</sup>, sea salt Cl<sup>-</sup>, sea salt Ca<sup>2+</sup>, sea salt and non-sea salt Mg<sup>2+</sup>, Na<sup>+</sup>, sea salt and non-sea salt K<sup>+</sup> indicating the natural source (sea and soil) contributions to rainwater. In addition, non-sea salt K<sup>+</sup> in rainwater presumably comes from biomass burning [1]. NH<sub>4</sub><sup>+</sup> sources might come from human and animal wastes and other human-made activities (possibly fertilizers, agriculture wastes, and urban activities).

Factor 2 explained approximately 10.78% of total variance, with strong positive loading for non-sea salt SO<sub>4</sub><sup>2-</sup>, NO<sub>3</sub><sup>-</sup>, H<sup>+</sup>, and non-sea salt Mg<sup>2+</sup> suggesting anthropogenic sources from vehicular emission, most notably fossil fuels combustion. With hydrogen ion being in the same factor, this also points to acidification as a result. In Bandung, there were ± 1.6 million units of a motor vehicle registered in 2015, which likely contributed significantly to SO<sub>2</sub> and NO<sub>3</sub> emissions [24].

Factor 3 explained approximately 8.76 % of the total variance with non-sea salt Cl<sup>-</sup> as the sole component, possibly as the result of the acid HCl with alkaline compound (Ca<sup>2+</sup>, Mg<sup>2+</sup>, and K<sup>+</sup>).

#### 4. CONCLUSIONS

The analysis of rainwater in Bandung in this study found that:

1. The pH of rainwater tends to be acidic with the values ranged between 3.4 to 7.71 (average 5.42 ± 0.72), with the average conductivity of 19.60 µS/cm.

2. The major anion component was SO<sub>4</sub><sup>2-</sup> (36.2% of the total anion mass), while NH<sub>4</sub><sup>+</sup> was the main cation component (47.7% of total cation mass). The major components found that, SO<sub>4</sub><sup>2-</sup> and NH<sub>4</sub><sup>+</sup> PWM concentrations were the highest at all sampling sites in almost every month, compared to other ions.
3. Ammonium (NH<sub>4</sub><sup>+</sup>) was found to be the dominant neutralizing ion.
4. Small-scale spatial variability found for nss-SO<sub>4</sub><sup>2-</sup>, nss-Cl<sup>-</sup>, nss-Ca<sup>2+</sup>, nss-Mg<sup>2+</sup>, and NH<sub>4</sub><sup>+</sup>, indicate the influence of local emission sources.
5. Principal Component Analysis (PCA) demonstrated that rainwater was also affected by natural sources of sea salts and crustal sources as well as biomass burning and local anthropogenic activities (vehicular emission, fossil fuels combustion, human and animal wastes and human made activities (fertilizers and agriculture wastes).

#### 5. ACKNOWLEDGMENTS

This paper is part of the work of the Riset KK dan Inovasi ITB 2017 entitled "Prediksi Dampak Kesehatan Lingkungan Deposisi Basah Logam Berat di Wilayah Perkotaan Bandung" funded by Institut Teknologi Bandung.

#### 6. REFERENCES

- [1] Sillapapiromsuk S. and Chantara S. Chemical Composition and Seasonal Variation of Acid Deposition in Chiang Mai, Thailand. Journal Environmental Engineering Resources, Vol. 15 (2), 2010, pp. 093-098
- [2] Budhavant K.B., Rao P.S.P., Safai P.D., and Ali, K. Chemistry of Monsoon and Post-Monsoon Rains at a High Altitude Location, Sinhagad, India. Journal Aerosol and Air Quality Research, Vol. 9(1), 2009, pp. 65 –79.
- [3] Wang H., and Han G. Chemical composition of rainwater and anthropogenic influences in Chengdu, Southwest China. Journal Atmospheric Research, Vol. 99, 2011, pp. 190-196.
- [4] Salve P.R., Gobre T., Lohkare H., Krupadam R.J., Bansawal A., Ramteke D.S., and Wate S.R. Source identification and variation in the chemical composition of rainwater at coastal and industrial areas of India. Journal Atmospheric Chemistry, Vol. 68, 2011, pp. 183-198.

- [5] Tiwari S., Chate D.M., Bisht D.S., Srivastava M.K., and Padmanabhamurty B. Rainwater chemistry in the North Western Himalayan Region, India. *Journal Atmospheric Research* Vol. 104-105, 2012, pp. 128-138.
- [6] Zhang X., Jiang H., Zhang Q., and Zhang X. Chemical characteristics of rainwater in northeast China, a case study of Dalian. *Atmospheric Research*, Vol.116, 2012, pp.151-160.
- [7] Cerón R.M., Cerón J.G., Carballo C.G., Aguilar C.A., Montalvo C., Benítez J.A., Y. J. Villareal Y.J. and Gómez M.M. Chemical Composition, Fluxes and Seasonal Variation of Acid Deposition in Carmen Island, Campeche, Mexico. *Journal of Environmental Protection*, Vol. 4, 2013, pp. 50-56.
- [8] Shrestha S., Pandey V.P., Yoneyama Y., Shrestha S. and Kazama F. An evaluation of rainwater quality in Kathmandu Valley, Nepal. *Journal Sustainable Environment Resources*, Vol. 23 (5), 2013, pp. 341-350.
- [9] Zhao M., Li L., Liu Z., Chen B., Huang J., Cai J., and Deng S. Chemical Composition and Sources of Rainwater Collected at a Semi-Rural Site in Ya'an, Southwestern China. *Atmospheric and Climate Sciences*, Vol. 3, 2013, pp. 486-496.
- [10] Gioda A., Olga L., Bracero M., Scatena F.N., Kathleen C., Weathers, Mateus V.L, and McDowell W.H. Chemical constituents in clouds and rainwater in the Puerto Rican rainforest: Potential sources and seasonal drivers. *Journal Atmospheric Environment*, Vol. 68, 2013, 208-220.
- [11] Cerqueira, F.R.F., Pinto M.F., Derossi, I.N., Esteves W.T., Santos M.D.R.S., and Matos R.C., Chemical characteristics of rainwater at a southeastern site of Brazil. *Atmospheric Pollution Research*, Vol. 5, 2014, pp. 253-261.
- [12] Cui J., Zhou J., Peng Y., He Y., Yang H., and Mao J, Zhang M., Wang Y. and Wang S. Atmospheric wet deposition of nitrogen and sulfur in the agroecosystem in developing and developed areas of Southeastern China. *Atmospheric Environment*, Vol. 89, 2014, pp. 102-108.
- [13] Niu H., He Y., Luc X.X, Shen J., Du J., Zhang T., Pu T., Xin H., and Chang L. Chemical composition of rainwater in the Yulong Snow Mountain region, Southwestern China. *Journal Atmospheric Research*, Vo. 144, 2014, pp. 195-206.
- [14] Pasha G.S.M., Shivashankara G.P., and Tiwari S. Chemical Composition and Ionic Variability of Wet and Bulk Precipitation in Urban Environment of Karnataka State, India, *International Journal Current Research and Academic Review*, Vol. 2(5), 2014, pp. 66 – 84.
- [15] Rai P. Wet Deposition of Gaseous Pollutant in Dhanbad India for Rain Water Quality. *International Journal of Current Engineering and Technology*, Vol. 4(2), 2014, pp. 673-679.
- [16] Zhang N., Cao J., He Y., and Xiao S. Chemical composition of rainwater at Lijiang on the Southeast Tibetan Plateau: influences from various air mass sources. *Journal Atmospheric Chemistry*, Vo. 51, 2014, pp.157-174.
- [17] Park S.M., Seo B.K., Lee G., Kahng S.H., and Jang Y.W. Chemical composition of water-soluble inorganic species in precipitation at Shinwa Basin, Korea, *Journal Atmosphere*, Vol. 6, 2015, pp. 732 – 75.
- [18] Tiwari S., Hopke P.K., Thimmaiah D., Dumka U.C., Srivastava A.K., Bisht D.S., Rao P.S.P., Chate D.M., Srivastava M.K., Tripathi S.N. Nature and Sources of Ionic Species in Precipitation across the Indo-Gangetic Plains, India. *Journal Aerosol and Air Quality Research*, Vol. 16, 2016, pp. 943-957.
- [19] Seinfeld, John, H., dan Pandis, S.N. *Atmospheric Chemistry and Physics from Air Pollution to Climate Change*, Second Edition, John Wiley and Son, Inc., New York. 2006.
- [20] Acid Deposition Monitoring Network (EANET), Technical Manual for Wet Deposition Monitoring in East Asia, Network Center for EANET, 2015.
- [21] Drijeana, Raper, D.W., and Gee, I.L. Spatial Variation and Sources Attribution of the Chemical Composition in Acid Deposition in the Peak District, Northern England. *Proceedings of The 95<sup>th</sup> Air & Waste Management Association Annual Conference & Exhibition*, Salt Lake City, UT., 2000.
- [22] Drijeana, Lee, D. S., Kingdon R. D., Raper, D. W. and Gee, I.L. (2001) : Modelling acid deposition in the UK at a scale of 5km by 5km, *Water, Air and Soil Pollution* 130, 319 – 324
- [23] Acid Deposition Monitoring Network (EANET). Third Periodic Report on the state of acid deposition in East Asia. Part I regional assessment, 2016.
- [24] BPS-Statistic of Bandung Municipality, 2016.



- [25] Cheng, Y., Liu, Y., Huo, M., Sun, Q., Wang, H., Chen, Z., and Bai. Chemical characteristics of precipitation at Nanping Mangdang Mountain in eastern China during spring. *Journal of Environmental Sciences*, Vol. 23(8), 2011, pp. 1350-1358.
- [26] Duan L., Yu Q., Zhang Q., Wang Z., Pan Y., Larssen T., Tang J. and Mulder J. Acid deposition in Asia: Emissions, deposition, and ecosystem effects. *Journal Atmospheric Environment*, Vol. 146, 2016, pp. 55-69.
- [27] World Meteorology Organization, Manual For The GAW Precipitation Chemistry Programme Guidelines, Data Quality Objectives and Standard Operating Procedures, GAW Precipitation Chemistry Science Advisory Group, 2004 Amended 2015.
- [28] Giannoni S.M., Rollenbeck R., Trachte K., and Bendix J. Natural or anthropogenic? On the origin of atmospheric sulfate deposition in the Andes of southeastern Ecuador. *Journal Atmospheric Chemistry and Physics*, Vol. 14, 2014, pp. 11297–11312.
- [29] Ianniello A., Spataro F., Esposito G., Allegrini I., Hu M., and Zhu T. Chemical characteristics of inorganic ammonium salts in PM<sub>2.5</sub> in the atmosphere of Beijing (China). *Journal Atmospheric Chemistry and Physics*, Vol. 11, 2011, pp. 10803–10822.
- [30] Xu Z., Wu Y., Liu W.J., Liang C.H., Ji, J., Zhao T., and Zhang X. Chemical composition of rainwater and the acid neutralizing effect at Beijing and Chizhou City. *Atmospheric Research*, Vol. 164-165, 2015, pp. 278-285.
- [31] Vet R., Artz R.S., Carou S., Shaw M., Ro C.U., Aas W., A., Baker, Bowersox V.C., Dentener F., Lacaux C.G., Hou A., Pienaar J.J., Gillett R., Forti C.M., Gromov S., Hara H., Khodzherm T., Mahowald N.M., Nickovic S., Rao P.S.P., and Reid N.W. A global assessment of precipitation chemistry and deposition of sulfur, nitrogen, sea salt, base cations, organic acids, acidity and pH, and phosphorus. *Journal Atmospheric Environment*, Vol. 93, 2014, pp. 3-100.
- [32] Xiao H.W., Xiao H.Y., Long A.M., Wang Y.L., and Liu C.Q. Chemical composition and source apportionment of rainwater at Guiyang, SW China. *Journal Atmospheric Chemistry*, Vol. 70, 2013, pp. 269-281.

---

Copyright © Int. J. of GEOMATE. All rights reserved, including the making of copies unless permission is obtained from the copyright proprietors.

---

# CHEMICAL COMPOSITION AND SOURCES ATTRIBUTION OF RAINWATER IN BANDUNG AREA, INDONESIA

## ORIGINALITY REPORT

19%

SIMILARITY INDEX

12%

INTERNET SOURCES

15%

PUBLICATIONS

3%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://www.mdpi.com">www.mdpi.com</a> Internet Source	1%
2	<a href="http://www.semanticscholar.org">www.semanticscholar.org</a> Internet Source	1%
3	<a href="http://businessdocbox.com">businessdocbox.com</a> Internet Source	1%
4	Tarigan, Ari K.M., Saut Sagala, D. Ary A. Samsura, Dika F. Fiisabilillah, Hendricus A. Simarmata, and Mangapul Nababan. "Bandung City, Indonesia", Cities, 2016. Publication	1%
5	<a href="http://www.nature.com">www.nature.com</a> Internet Source	1%
6	Wu, Qixin, and Guilin Han. "Sulfur isotope and chemical composition of the rainwater at the Three Gorges Reservoir", Atmospheric Research, 2015. Publication	1%
7	<a href="http://res.mdpi.com">res.mdpi.com</a>	

Internet Source

1 %

8

Acid rain 2000, 2001.

Publication

1 %

9

K.B. Budhavant, P.S.P. Rao, P.D. Safai, K. Ali.

"Influence of local sources on rainwater chemistry over Pune region, India",

Atmospheric Research, 2011

Publication

1 %

10

Submitted to University College London

Student Paper

1 %

11

etd.aau.edu.et

Internet Source

1 %

12

iopscience.iop.org

Internet Source

<1 %

13

pea.lib.pte.hu

Internet Source

<1 %

14

Yu Zhao, Qi Feng, Aigang Lu. "Variation of chemical characteristics of precipitation with respect to altitude gradient on the northern slope of Mt. Taibai, China", Journal of Atmospheric and Solar-Terrestrial Physics, 2020

Publication

<1 %

15

Jianping Wu, Yongxian Su, Xiuzhi Chen, Liyang Liu et al. "Redistribution characteristics of

<1 %

atmospheric precipitation in different spatial levels of Guangzhou urban typical forests in southern China", Atmospheric Pollution Research, 2019

Publication

---

16

Zhao, Min, Li Li, Zhilin Liu, Bin Chen, Jianqiu Huang, Jinwang Cai, and Shihuai Deng. "Chemical Composition and Sources of Rainwater Collected at a Semi-Rural Site in Ya'an, Southwestern China", Atmospheric and Climate Sciences, 2013.

Publication

---

17

Qixin Wu, Guilin Han. "Sulfur isotope and chemical composition of the rainwater at the Three Gorges Reservoir", Atmospheric Research, 2015

Publication

---

18

Rao, P.S.P., S. Tiwari, J.L. Matwale, S. Pervez, P. Tunved, P.D. Safai, A.K. Srivastava, D.S. Bisht, S. Singh, and P.K. Hopke. "Sources of chemical species in rainwater during monsoon and non-monsoonal periods over two mega cities in India and dominant source region of secondary aerosols", Atmospheric Environment, 2016.

Publication

---

19

Weiwei Pu, Weijun Quan, Zhiqiang Ma, Xuefeng Shi, Xiujuan Zhao, Linna Zhang,

<1 %

<1 %

<1 %

<1 %

Zhenfa Wang, Wenyan Wang. "Long-term trend of chemical composition of atmospheric precipitation at a regional background station in Northern China", Science of The Total Environment, 2017

Publication

20

[online.ucpress.edu](http://online.ucpress.edu)

Internet Source

<1 %

21

Submitted to Asian Institute of Technology

Student Paper

<1 %

22

Sase, H.. "Seasonal variation in the atmospheric deposition of inorganic constituents and canopy interactions in a Japanese cedar forest", Environmental Pollution, 200803

Publication

<1 %

23

[downloads.hindawi.com](http://downloads.hindawi.com)

Internet Source

<1 %

24

[www.chem.itb.ac.id](http://www.chem.itb.ac.id)

Internet Source

<1 %

25

A.I. Calvo, V. Pont, F.J. Olmo, A. Castro, L. Alados-Arboledas, A.M. Vicente, M. Fernández-Raga, R. Fraile. "Air Masses and Weather Types: A Useful Tool for Characterizing Precipitation Chemistry and Wet Deposition", Aerosol and Air Quality Research, 2012

<1 %

26

S. Sillapapiromsuk. "Chemical Composition and Seasonal Variation of Acid Deposition in Chiang Mai, Thailand", Environmental Engineering Research, 06/30/2010

Publication

---

<1 %

27

confbeam.org

Internet Source

---

<1 %

28

Hatice Karadeniz, Serpil Yenisoy-Karakaş. "Evaluation of hourly-based precipitation chemistry in suburban site of Bolu", Atmospheric Pollution Research, 2020

Publication

---

<1 %

29

Qiangmei Leng, Jian Cui, Fengwu Zhou, Ke Du et al. "Wet-only deposition of atmospheric inorganic nitrogen and associated isotopic characteristics in a typical mountain area, southwestern China", Science of The Total Environment, 2018

Publication

---

<1 %

30

Wen Xu, Zhang Wen, Bo Shang, Anthony J. Dore et al. "Precipitation chemistry and atmospheric nitrogen deposition at a rural site in Beijing, China", Atmospheric Environment, 2020

Publication

---

<1 %

31

www.environmentportal.in

Internet Source

<1 %

32

Jie Zeng, Guilin Han. "Rainwater Chemistry Reveals Air Pollution in a Karst Forest: Temporal Variations, Source Apportionment, and Implications for the Forest", Atmosphere, 2020

Publication

&lt;1 %

33

Marcos Rodrigues Facchini Cerqueira, Marcelo Fonseca Pinto, Ingrid Nunes Derossi, Wesley Tinoco Esteves et al. "Chemical characteristics of rainwater at a southeastern site of Brazil", Atmospheric Pollution Research, 2014

Publication

&lt;1 %

34

Monika J. Kulshrestha, Ruchi Singh, R. Duarah, P.G. Rao. "Influence of crustal aerosols on wet deposition at a rural site of North-East India", International Journal of Environmental Studies, 2014

Publication

&lt;1 %

35

Submitted to Universitas Pelita Harapan

Student Paper

&lt;1 %

36

Zhifang Xu, Yao Wu, Wen-Jing Liu, Chong-Shan Liang, Jianpeng Ji, Tong Zhao, Xuan Zhang. "Chemical composition of rainwater and the acid neutralizing effect at Beijing and Chizhou city, China", Atmospheric Research, 2015

Publication

&lt;1 %

37

[cdrjournal.com](http://cdrjournal.com)

Internet Source

<1 %

38

Ágnes Keresztesi, Marius-Victor Birsan, Ion-Andrei Nita, Zsolt Bodor, Róbert Szép.

"Assessing the neutralisation, wet deposition and source contributions of the precipitation chemistry over Europe during 2000–2017", Environmental Sciences Europe, 2019

Publication

<1 %

39

Eduardo Henrique Martins, Danilo Covaes Nogarotto, Jefferson Mortatti, Simone Andréa Pozza. "Chemical Composition of Rainwater in an Urban Area of the Southeast of Brazil", Atmospheric Pollution Research, 2018

Publication

<1 %

40

Guozheng Zhang, Duanyang Liu, Xiaohong He, Deyue Yu, Meijuan Pu. "Acid rain in Jiangsu province, eastern China: Tempo-spatial variations features and analysis", Atmospheric Pollution Research, 2017

Publication

<1 %

41


Lin Ma, Hossein Dadashazar, Miguel Ricardo A. Hilario, Maria Obiminda Cambaliza et al. "Contrasting wet deposition composition between three diverse islands and coastal North American sites", Atmospheric Environment, 2020

Publication

<1 %



42

Pascaud, A., S. Sauvage, P. Coddeville, M. Nicolas, L. Crois , A. Mezdour, and A. Probst. "Contrasted spatial and long-term trends in precipitation chemistry and deposition fluxes at rural stations in France", Atmospheric Environment, 2016.

Publication

&lt;1 %

43

asianjae.org

Internet Source

&lt;1 %

44

mafiadoc.com

Internet Source

&lt;1 %

45

Guilin Han, Qixin Wu, Yang Tang. "Acid rain and alkalization in southwestern China: chemical and strontium isotope evidence in rainwater from Guiyang", Journal of Atmospheric Chemistry, 2012

Publication

&lt;1 %

46

Hung-Yu Chen, Ling-Fei Hsu, Shih-Zhe Huang, Long Zheng. "Assessment of the Components and Sources of Acid Deposition in Northeast Asia: A Case Study of the Coastal and Metropolitan Cities in Northern Taiwan", Atmosphere, 2020

Publication

&lt;1 %

47

Jing Li, Huawu Wu, Pengju Jiang, Congsheng Fu. "Rainwater chemistry in a subtropical high-altitude mountain site, South China:

&lt;1 %

Seasonality, source apportionment and potential factors", Atmospheric Environment, 2021

Publication

48

Md Firoz Khan, Khairul Nizam Abdul Maulud, Mohd Talib Latif, Jing Xiang Chung et al. "Physicochemical factors and their potential sources inferred from long-term rainfall measurements at an urban and a remote rural site in tropical areas", Science of The Total Environment, 2018

Publication

<1 %

49

Sakihama, H.. "Chemical characteristics of precipitation in Okinawa Island, Japan", Atmospheric Environment, 200803

Publication

<1 %

50

[hdl.handle.net](https://hdl.handle.net)

Internet Source

<1 %

Exclude quotes Off

Exclude matches Off

Exclude bibliography On