

**POLYTECHNIC HEALTH KEMENKES BANDUNG MIDWIFERY STUDY
PROGRAM BOGOR FINAL PROJECT REPORT, JUNE. 2024**

Dwi Nur Anggraini

P17324221053

**Pregnancy Obstetrics Care for Mrs. D 18 Years Old G1P0A0 31 Weeks Pregnancy
Age with Condyloma Acuminata**

VI CHAPTER, 99 pages, 8 appendix, 4 figures

ABSTRACT

Pregnancy at the age of 18 years is a teenage pregnancy that can cause several risks to both the mother and the fetus, for the fetus there can be complications of premature birth and LBW. In mothers, bleeding, birth canal tears, and prolonged labor can occur. In 2018 as many as 12.8 million mothers gave birth aged 15-19 years. In this case, the mother had a young pregnancy at the age of 18 accompanied by condyloma acuminata, which has the potential to cause respiratory problems, vocal cord damage, and eye infections in HPV-infected babies.

The method used in the preparation of this Final Project Report is a case report, midwifery management approach and SOAP documentation (Subjective, Objective, Analysis, Management). Data collection techniques use interview techniques, physical examination, observation, documentation studies, and literature studies.

Subjective data results on Mrs. D, age 18 years, HPHT (First Period Last Day) August 21, 2023, TP (Interpretation of Labor) May 28, 2024, gestational age 31 weeks, is the first child, has never miscarried, there is a lump in the vagina. Objective data from abdominal examination was palpable round, hard, bouncy (head), hard, flat like a board on the right side (puka), round, soft (buttocks). On genital examination there are no varicose veins, no Bartholin and Skene glands, there are lumps resembling cauliflowers like cockscombs on the right and left labia minora. After a supporting examination, the results of laboratory tests for syphilis, HIV / AIDS, and HBsAg were non-reactive. Analysis was established that Mrs. D, age 18 years G1P0A0, gestational age 31 weeks with condyloma acuminata single fetus living intrauterine. Management carried out in collaboration with a dermatologist for the administration of Trichloroacetic acid 80 - 90% and gentamicin sulfate ointment 0.1% to be applied to condyloma acuminata, while for the management of adolescent pregnancy, pregnancy and childbirth counseling is carried out.

The conclusion of the results of care is in accordance with the standard of care for pregnancy services, but there are obstacles, namely an uncooperative husband. Advice given to Mrs. D invites her husband for treatment so that sexually transmitted diseases in this couple are resolved.

Keywords : Obstetric care of pregnancy, Condyloma Acuminata

Bibliography : 30 (2018-2024)