

# THE EFFECT OF DIFFERENCES YEAST'S CONCENTRATION ON TOTAL ANTHOCYANIN CONTENT, AND ANTIOXIDANT ACTIVITIES OF BLACK TAPE EXTRACT (ORYZA SATIVA L VAR FORMA GLUTINOSA)

Trinovani, Elvi<sup>1</sup>, Wahdah, Hanifah Nur Fauziyah<sup>1</sup>, Fauziyah, Roro Nur<sup>2</sup>

Department of Pharmacy Health Polytechnic Ministry of Health, Bandung, Indonesia<sup>1</sup>  
Department of Nutrition Health Polytechnic Ministry of Health, Bandung, Indonesia<sup>2</sup>



**ABSTRACT**— Anthocyanin is a sub-type organic substance from the flavonoid, and is one example of natural- colored pigments. This substance can be found in food items such as Fermented Black Glutinous Rice. Aside from anthocyanin, previously studied Fermented Black Glutinous Rice samples also possess antioxidant activity that can be developed to prevent hyperlipidemia. However, the fermenting process on the black rice is known to affect the anthocyanin substance and the antioxidant activity contained within the black rice. This study is conducted to find out the specific effects caused by the yeast and its concentrations on the fermenting process. The results show that the yeast concentrations do affect the anthocyanin level and the antioxidant activity in the black rice. The anthocyanin level, after subjected to some pH differential tests, plummeted as the level of yeast concentrations went up. The highest anthocyanin level was obtained in the sample with 0.5% yeast concentration which was equal to 566,0226 mg / 100g. Likewise, the results of examination of antioxidant activity have decreased with increasing yeast concentration. The highest antioxidant activity was obtained at 0.5% yeast concentration. This is showed by the IC<sub>50</sub> value of 81.9834 ppm which means the sample has strong antioxidant activity.

**KEYWORDS:** Fermented Black Glutinous Rice, Anthocyanin, Antioxidant Activity.

## 1. INTRODUCTION

Anthocyanins are a sub-type of organic compounds from the flavonoid family. Some of the most common anthocyanin compounds are pelargonidin, peonidin, cyanidin, malvidin, petunidin, and delphinidin [23]. Anthocyanins work as secondary antioxidants as is the case with  $\beta$ -carotene, which breaks down the oxidacidipid peroxide chain. Antioxidants are free radical scavengers or antidotes [27]. The source of anthocyanin and fiber besides fruits and vegetables is rice (*Oryza sativa*) which is rich in anthocyanin such as black Glutinous Black Rice, black rice and brown rice [18]. Fermented Glutinous Black Rice (*Oryza sativa glutinosa*) as a raw material for Fermented Glutinous Black Rice tape is a very potential commodity as a source of carbohydrates, antioxidants, bioactive compounds and fibers that are important for health [38]. One of the foods in Indonesia made from Glutinous Black Rice is Fermented Glutinous Black Rice containing ferocyanin, phenols and antioxidant activity. Glutinous tape has anthocyanin content of 257 ppm or equivalent to 25.7 mg / 100 grams. [8]. Glutinous Black Rice tape is a traditional Indonesian food made from Glutinous Black Rice which is processed by fermentation. In the Glutinous Black Rice fermentation process the preparation (washing, cooking and cooling) stages, workmanship, and fermentation (heating, inoculation with yeast, and incubation) are important influential stages to produce good quality Glutinous Black Rice tape [16]. The process of making Fermented Glutinous Black Rice is done by washing Fermented Glutinous Black Rice to clean dirt and contamination. Furthermore, black glutinous rice soaked for several hours. Soaking aims to

produce Fermented Glutinous Black Rice that is not hard and shorten the steaming time. Then black rice that has been soaked, boiled half done and then steamed [9]. The steaming process will cause the starch to gelatinize and will break into amylose and amylopectin. Gelatinized starch is used as a growth medium for microbes present in yeast. Before the fermentation of Glutinous Black Rice is carried out, it is cooled first until the temperature approaches room temperature so that the microbes present in the yeast can work optimally. Add yeast with a concentration of 0.1% -0.5% because at that concentration will produce a tape with a sweet taste, sour and distinctive aroma tape. The yeast used for fermentation is *Sacharomyces cereviseae* [9].

In addition to influencing the taste of food, the provision of yeast in the fermentation process can affect the ingredients in food. Among them the fermentation method which gives a very significant effect on anthocyanin levels of sweet potato flour during the fermentation process [22]. It is also as stated by Medina (2018) that yeast has been shown to interact with anthocyanins for example is interactions with yeast cell wall material. So that it allows a decrease in anthocyanin levels during the Fermented Glutinous Black Rice fermentation process. The use of varied yeast in each yeast-producing region can also be one of the factors that influence the fermentation process. Cililin area, West Bandung Regency is one of the regions producing Fermented Glutinous Black Rice. The use of yeast during Fermented Glutinous Black Rice fermentation in the Cililin area still uses concentrations in accordance with a pre-existing recipe of 0.1%. At the concentration of the yeast indeed obtained tape with optimum conditions. However, there is no known concentration of yeast that can provide maximum levels of anthocyanin and antioxidant activity on Fermented Glutinous Black Rice. Based on the background that has been explained, it is known that there are many benefits of anthocyanin, but there are several things that can affect anthocyanin levels when making Fermented Glutinous Black Rice tape and the need to illustrate the effect of yeast concentration on anthocyanin levels and antioxidant activity of Fermented Glutinous Black Rice in order to obtain optimum yeast concentration. So the authors are very interested in conducting a study entitled "Effect of Differences in Yeast Concentration on Total Anthocyanin Levels, And Antioxidant Activity of Fermented Glutinous Black Rice Extract (*Oryza sativa* L var forma glutinosa).

## **2. Method**

This type of research is a descriptive study to determine the effect of different levels of yeast given on total anthocyanin levels and antioxidant activity on Fermented Glutinous Black Rice. This study uses Fermented Glutinous Black Rice obtained from Cililin in West Bandung Regency, West Java. The study was conducted at the Bandung Health Polytechnic Laboratory, Department of Pharmacy. The time of the research was carried out in March 2019. Materials used included test materials and chemicals. Test materials used are Black Glutinous Rice and Chemicals used include Aquabides, KCl 0.2 N solution; HCl 0.2 N solution; potassium acetate, ascorbic acid, methanol pa merck brand, DPPH (1,1-diphenyl-2-picrylhydrazyl), [(CH<sub>3</sub>COO) 2Cu.H<sub>2</sub>O], [(CH<sub>3</sub>COO) 2Mg.4H<sub>2</sub>O], ethanol pa brand merck, filter paper whatman no. 42, aluminum foil and tissue roll. The tools used include UV-Vis spectrophotometers, glass cuvettes, pH meters, analytical scales, water baths, centrifugers, mercury thermometers, glassware, stopwatches, filters and parchment paper. The workings of this study began with the preparation of test material in the form of Fermented Glutinous Black Rice. Then the fermentation process is carried out, the results of which will be tested, namely, identification of anthocyanin pigment compounds, analysis of maximum wavelength, determination of anthocyanin levels and antioxidant activity test using DPPH method

## **3. Result**

The plants used in this research are Fermented Glutinous Black Rice (*Oryza sativa* L.) Poaceae tribe which has been fermented into Fermented Glutinous Black Rice taken from the Poltekkes Bandung Village of Cililin

Village, West Bandung Regency. This research was conducted using 5 groups of Fermented Glutinous Black Rice samples with several different concentrations of yeast. Before testing the sample, extraction is done first with the aim of being able to take anthocyanin compounds that are deposited in the sample. In this research, extraction was done using maceration method which was then continued with freeze-dry. The sample extraction results obtained are in the table.

**Table 1** Yield of Fermented Glutinous Black Rice Extract with 0.05% Yeast Concentration

Solvent	Methanol + HCl 1%
Sample Weight	250 g
Extract Weight	20,8494 g
Yield Extract	8,34 %

**Table 2** Yield of Fermented Glutinous Black Rice Extract with 0.075% Yeast Concentration

Solvent	Methanol + HCl 1%
Sample Weight	250 g
Extract Weight	23,7579 g
Yield Extract	9,50 %

**Table 3** Yield of Fermented Glutinous Black Rice Extract with 0.1% Yeast Concentration

Solvent	Methanol + HCl 1%
Sample Weight	250 g
Extract Weight	21,1781 g
Yield Extract	8,54 %

**Table 4** Yield of Fermented Glutinous Black Rice Extract with 0.125% Yeast Concentration

Solvent	Methanol + HCl 1%
Sample Weight	250 g
Extract Weight	22,3693 g
Yield Extract	8,95 %

**Table 5** Yield of Fermented Glutinous Black Rice Extract with 0.15% Yeast Concentration

Solvent	Methanol + HCl 1%
Sample Weight	250 g
Extract Weight	19,6600 g
Yield Extract	7,86 %

Anthocyanin identification can be done by doing color reagents by reacting the extract with HCl and NaOH. The results of identifying the color of Fermented Glutinous Black Rice extract with color reagents are in table 6.

**Table 6** Results of Anthocyanin Identification with Color Reagents

Reactor	Color	Identification results
HCl	Purplish red	+
NaOH	Color is fading	+

Determination of anthocyanin levels is done by the pH differential method. Determination of anthocyanin levels is done by using a pH difference that is pH 1.0 and pH 4.5. This is done because at pH 1.0 anthocyanin forms a red flavylum / oxonium kation. Meanwhile, at a pH of 4.5 anthocyanins form a colorless carbinol compound. Thus, the determination of levels can be done by making aliquots of anthocyanin solutions in pH 1.0 and pH 4.5 to then be measured for absorbance. The sample used was Fermented Glutinous Black Rice extract with some concentration of yeast. Samples that have been dissolved with different pH buffer solutions are then inserted in a spectrophotometer to determine the absorbance of each solution. Following is the absorbance results table for each Fermented Glutinous Black Rice extract solution with a yeast concentration of 0.05%.

**Table 7** Anthocyanin levels of Fermented Glutinous Black Rice Extract 0.05% Yeast Concentration

	Replication	1	2	3
pH 4,5	515 nm	0,465	0,477	0,484
	700 nm	0,200	0,219	0,204
pH 1	513 nm	2,356	2,263	2,216
	700 nm	0,644	0,600	0,604
Anthocyanin Absorption		1,447	1,405	1,332
Anthocyanin levels (mg/100g)		604,0822	586,5483	556,0729
Average of Anthocyanin levels		582,2344 mg/100g		

Absorbance samples that have been measured using a spectrophotometer in wavelengths of 510 nm and 700 nm can be calculated for anthocyanin levels by determining the absorbance of the sample first. The absorbance of each solution is calculated based on the equation found in the example calculation. The total anthocyanin monomeric from the dried extract of Fermented Glutinous Black Rice was calculated as cyaniding-3glucoside. The results of anthocyanin levels from methanol extract and HCl Fermented Glutinous Black Rice tape with a concentration of 0.75% can be seen in table 8

**Table 8** Anthocyanin Levels of Fermented Glutinous Black Rice Extract 0.075% Yeast Concentration

	Replication	1	2	3
pH 4,5	515 nm	0,328	0,329	0,331
	700 nm	0,177	0,173	0,181
pH 1,0	513 nm	1,919	1,987	1,922
	700 nm	0,533	0,523	0,525
Anthocyanin Absorption		1,303	1,308	1,247
Anthocyanin levels (mg/100g)		543,9662	546,0535	520,5877
Average of Anthocyanin levels		536,8691mg/100g		

From the table calculation of anthocyanin levels above, obtained anthocyanin levels in Fermented Glutinous Black Rice extract with a yeast concentration of 0.5% amounted to 536.8691 mg / 100g. The results of anthocyanin levels from methanol extract and HCl Fermented Glutinous Black Rice with a concentration of 1% can be seen in table 9

**Table 9** Anthocyanin Levels of Fermented Glutinous Black Rice Extract 0.1% Yeast Concentration

	Replication	1	2	3
pH 4,5	515 nm	0,283	0,268	0,231
	700 nm	0,106	0,100	0,094
pH 1,0	513 nm	1,364	1,354	1,295
	700 nm	0,278	0,304	0,283
Anthocyanin Absorption		0,909	0,882	0,875
Anthocyanin levels (mg/100g)		379,4822	368,2104	365,2881
Average of Anthocyanin levels		370,9936 mg/100g		

From the calculation table of anthocyanin levels above, obtained anthocyanin levels in Fermented Glutinous Black Rice extract with 1% yeast concentration is 370.9936 mg / 100g. The results of anthocyanin test results from methanol extract and HCl Fermented Glutinous Black Rice with a concentration of 1.25% can be seen in table 10.

**Table 10** Anthocyanin levels of Fermented Glutinous Black Rice Extract Yeast Concentration 0.125%

	Replication	1	2	3
pH 4,5	515 nm	0,127	0,111	0,113
	700 nm	0,092	0,089	0,089
pH 1,0	513 nm	0,663	0,666	0,692
	700 nm	0,155	0,166	0,189
Anthocyanin Absorption		0,473	0,478	0,479
Anthocyanin levels (mg/100g)		197,4643	199,5517	199,9691
Average of Anthocyanin levels		198,9950 (mg/100g)		

From the calculation of anthocyanin levels above, obtained anthocyanin levels in Fermented Glutinous Black Rice extract with a yeast concentration of 1.25% amounted to 198.995 mg / 100g. The results of anthocyanin test results from methanol extract and HCl Fermented Glutinous Black Rice with a concentration of 0.15% can be seen in table 11. From the calculation table above, obtained anthocyanin levels in Fermented Glutinous Black Rice extract with a yeast concentration of 0.15% amounted to 189.1149 mg / 100g.

**Table 11** Anthocyanin levels of Fermented Glutinous Black Rice Extract Yeast Concentration of 0.15%

	Replication	1	2	3
pH 1,0	515 nm	0,104	0,106	0,105
	700 nm	0,083	0,087	0,088

pH 4,5	513 nm	0,544	0,540	0,543
	700 nm	0,064	0,071	0,076
Anthocyanin Absorption		0,459	0,450	0,450
Anthocyanin levels (mg/100g)		191,6197	187,8625	187,8625
Average of Anthocyanin levels			189,1149 (mg/100g)	

From all samples of Fermented Glutinous Black Rice extract that have been tested, there are differences in anthocyanin levels. The results of the calculation of anthocyanin levels from Fermented Glutinous Black Rice extract with various yeast concentrations can be seen in table 12.

**Table 12** Results of Anthocyanin Analysis on Fermented Glutinous Black Rice Extract

Sample	pH 4.5		pH 1.0		Anthocyanin Absorption	Anthocyanin levels (mg/100g)
	515 nm	700 nm	515 nm	700 nm		
Sample <sup>1</sup>	0,475	0,208	2,240	0,616	1,356	566,0226
Sample <sup>2</sup>	0,329	0,177	1,955	0,527	1,275	532,3465
Sample <sup>3</sup>	0,261	0,100	1,338	0,288	0,889	370,9936
Sample <sup>4</sup>	0,117	0,090	0,674	0,170	0,477	198,995
Sample <sup>5</sup>	0,105	0,086	0,542	0,070	0,453	189,1149

Note:

- 1 Samples with 0.05% yeast concentration
- 2 Samples with 0.075% yeast concentration
- 3 Samples with 0.1% yeast concentration
- 4 Samples with 0.125% yeast concentration
- 5 Samples with 0.15% yeast concentration


One method that is often used to determine the antioxidant activity of Fermented Glutinous Black Rice is the DPPH method. In measuring antioxidant activity, five concentrations of using yeast during fermentation were used to determine the effect of yeast during fermentation on the antioxidant activity of Fermented Glutinous Black Rice. The antioxidant activity of black crab tape with some yeast concentration during fermentation is indicated by the IC<sub>50</sub> value. The results of antioxidant activity tests of methanol extract and HCl Fermented Glutinous Black Rice with a concentration of 0.05% can be seen in table 13. The data in table 13 is the absorbance data of Fermented Glutinous Black Rice extract with a yeast concentration of 0.05% obtained after measurements using a UV-Vis spectrophotometer. Obtained a decrease in absorbance along with an increase in the concentration of the sample solution. Absorbance data is then calculated% inhibition or% of antioxidant activity with a predetermined formula. Then the results are used to calculate

**Table 13** Antioxidant Activity of Fermented Glutinous Black Rice Extract 0.05% Yeast Concentration

Concentration (ppm)	Absorbance	% Inhibition	Equation (y=bx+c)	IC <sub>50</sub> (ppm)
0	0,985	0	y = 0,3856x +	81,983

20	0,574	41,72589	18,3872 $R^2 = 0,5935$
40	0,5663	42,50761	
60	0,5483	44,33503	
80	0,5227	46,93401	
100	0,4877	50,48731	

IC<sub>50</sub> which is the value to measure antioxidant activity with the linear equation  $y = ax + b$ . IC<sub>50</sub> equation that has been obtained can be illustrated by graph 1.




**Graph 1** Antioxidant Activity of Fermented Glutinous Black Rice Extract 0.05% Yeast Concentration

The results of antioxidant activity tests of methanol extract and HCl Fermented Glutinous Black Rice with a concentration of 0.075% can be seen in table 14. Based on table 14 IC<sub>50</sub> values or the concentration of methanol extract and HCl Fermented Glutinous Black Rice tape with a yeast concentration of 0.075% inhibiting 50% of DPPH free radicals at 100.531 ppm.

**Table 14** Antioxidant Activity of Fermented Glutinous Black Rice Extract 0.075% Yeast Concentration

Concentration (ppm)	Absorbance	% Inhibition	Equation (y=bx+c)	IC <sub>50</sub> (ppm)
0	0,985	0	$y = 0,3207x + 17,7597$ $R^2 = 0,5208$	100,531
20	0,6083	38,24365		
40	0,5957	39,52284		
60	0,5847	40,63959		
80	0,5727	41,85787		
100	0,5663	42,50761		

IC<sub>50</sub> equation that has been obtained can be illustrated by graph 2.



**Graph 2** Antioxidant Activity of Fermented Glutinous Black Rice Extract 0.075% Yeast Concentration


The results of antioxidant activity tests of methanol extract and HCl Fermented Glutinous Black Rice with a concentration of 0.1% can be seen in table 15.

**Table 15** Antioxidant Activity of Fermented Glutinous Black Rice Extract Yeast Concentration 0.1%

Concentration (ppm)	Absorbance	% Inhibition	Equation (y=bx+c)	IC50 (ppm)
0	0,985	0	y = 0,3313x + 15,9990	102,629
20	0,6397	35,05584	R <sup>2</sup> = 0,58806	
40	0,6163	37,43147		
60	0,597	39,39086		
80	0,5793	41,18782		
100	0,5683	42,30457		

Based on table 15, the IC50 value or the concentration of methanol extract and HCl Fermented Glutinous Black Rice tape with a yeast concentration of 0.1% inhibiting 50% of DPPH free radicals is 102.662 ppm.

IC50 equation that has been obtained can be illustrated by graph 3.



**Graph 3** Antioxidant Activity of Fermented Glutinous Black Rice Extract Yeast Concentration 0.1%

The results of antioxidant activity tests of methanol extract and HCl Fermented Glutinous Black Rice with a




concentration of 0.125% can be seen in table 16.

**Table 16.** Antioxidant Activity of Fermented Glutinous Black Rice Extract Yeast Concentration 0.125%

Concentration (ppm)	Absorbance	% Inhibition	Equation (y=bx+c)	IC50 (ppm)
<b>0</b>	0,985	0	y = 0,3241x + 14,6343 R <sup>2</sup> = 0,6191	109,12
<b>20</b>	0,654	33,6041		
<b>40</b>	0,643	34,7208		
<b>60</b>	0,6327	35,7665		
<b>80</b>	0,6003	39,0558		
<b>100</b>	0,5723	41,8985		

Based on table 4.15, the IC50 value or the concentration of methanol extract and HCl Fermented Glutinous Black Rice tape with a yeast concentration of 0.125% inhibits 50% of DPPH free radicals at 109.12 ppm.

IC50 equation that has been obtained can be illustrated by graph 4.



**Graph 4.** Antioxidant Activity of Fermented Glutinous Black Rice Extract Yeast Concentration 0.125%


The results of antioxidant activity tests of methanol extract and HCl Fermented Glutinous Black Rice with a concentration of 0.15% can be seen in table 17

**Table 17** Antioxidant Activity of Fermented Glutinous Black Rice Extract Yeast Concentration of 0.15%

Concentration (ppm)	Absorbance	% Inhibition	Equation (y=bx+c)	IC50 (ppm)
<b>0</b>	0,985	0	y = 0,3542x + 9,5915 R <sup>2</sup> = 0,81913	114,084
<b>20</b>	0,75	23,858		
<b>40</b>	0,7007	28,863		
<b>60</b>	0,6567	33,330		

<b>80</b>	0,6217	36,883
<b>100</b>	0,5823	40,883

Based on table 17, the IC50 value or the concentration of methanol extract and HCl Fermented Glutinous Black Rice tape with a yeast concentration of 0.15% inhibits 50% of DPPH free radicals of 114.08192 ppm. IC50 equation that has been obtained can be illustrated by graph 5.



**Graph 5** Antioxidant Activity of Fermented Glutinous Black Rice Extract Yeast Concentration of 0.15%

From all the Fermented Glutinous Black Rice extract samples that were tested there were differences in IC50 values. IC50 calculation results from Fermented Glutinous Black Rice extract with various yeast concentrations can be seen in table 18.

**Table 18** Results of Antioxidant Activity Analysis of Fermented Glutinous Black Rice Extract


<b>Sample</b>	<b>IC 50 (ppm)</b>
<b>1</b>	81.9834
<b>2</b>	100.531
<b>3</b>	102.629
<b>4</b>	109.12
<b>5</b>	114,084

Note:

- 1 Samples with 0.05% yeast concentration
- 2 Samples with 0.075% yeast concentration
- 3 Samples with 0.1% yeast concentration
- 4 Samples with 0.125% yeast concentration
- 5 Samples with 0.15% yeast concentration

Based on table 18 IC50 values or the concentration of methanol extract and HCl Fermented Glutinous Black Rice tape with some yeast concentrations inhibiting 50% of DPPH free radicals increased with an increase in yeast concentration. IC7 reduction that has been obtained can be illustrated by graph 6. The comparison used in this study was quercetin which was also tested by DPPH by measuring the absorbance of the comparator after adding DPPH reagents first. Then, the comparative absorbance results are used to calculate the 50%

inhibitory concentration of free radicals / IC50. The results of antioxidant activity tests from quercetin comparator can be seen in table 19




**Graph 6** Comparison of Antioxidant Activity for each Concentration

**Table 19.** Antioxidant Activity of Quercetin

Concentration (ppm)	Absorbance	% Inhibition	Equation (y=bx+c)	IC50 (ppm)
0	0,774	0	$y = 3,7652x + 12,668$	9,915
5	0,53	31,5245	$R^2 = 0,8995$	
10	0,298	61,4987		
15	0,135	82,5581		
20	0,071	90,8269		
25	0,062	91,9897		

Based on table 19, the IC50 value or quercetin comparative concentration inhibiting 50% of DPPH free radicals is 9,915 ppm. IC50 equation that has been obtained can be illustrated by graph 7.



**Graph 7** Quercetin Comparative Antioxidant Activity

#### 4. Discussion

This research was carried out on Fermented Glutinous Black Rice with five concentrations of yeast

fermentation to determine its effect on anthocyanin levels and antioxidant activity. Before testing the sample, extraction is done first. Extraction is the first step in determining the characterization of pigments so that coarse extracts are obtained. In fruits or vegetables, anthocyanin pigments are generally found in cells located near the surface. Anthocyanin contained in these tissues can be obtained by extraction using certain solvents. One theory says that the extracting agent can cause denaturation of the cell membrane so that the pigment contained in the membrane can be extracted [9]. The effectiveness of the extraction process is inseparable from the ability of the extracting material to dissolve the extracted component. Dissolution event of a substance occurs because of the interaction between the solvent and the dissolved material and can be divided into three stages, namely, the stage of breaking bonds between molecules of solutes that require energy, the stage of breaking bonds between solvent molecules that require energy, and finally the stage of formation a bond between a solute molecule and an energy-solvent molecule. If the energy produced is greater than the energy required, the dissolution process will occur. Polar compounds can only dissolve in polar solvents, as well as non-polar compounds which can only be dissolved in non-polar solvents as well [34]. This maceration process is carried out twice 24 hours to produce a purple filtrate. This is done to optimize the extraction process so that the anthocyanin pigment contained in the Fermented Glutinous Black Rice can be extracted entirely. Stirring with the stirrer is done to increase the effectiveness of the extraction process. The resulting filtrate is then concentrated by the freeze-dry method. The use of low temperatures aims to avoid the degradation and hydrolysis of anthocyanin pigments (Timberlake and Bridle, 1983). The extract powder obtained is then put in a dark bottle and stored in the freezer. Storage in the freezer aims to maintain the stability of anthocyanin which is very easily degraded. This analysis was carried out using a UVV is spectrophotometer at two wavelengths, 515 and 700 nm. The wavelength of 515 nm is the maximum wavelength of anthocyanin Fermented Glutinous Black Rice. This result is obtained by dissolving the Fermented Glutinous Black Rice extract in buffer pH 1 which is then measured its maximum wavelength.

The average results of anthocyanin levels from Fermented Glutinous Black Rice extract with 0.05% yeast concentration was 566.0226 mg / 100g, for Fermented Glutinous Black Rice extract with 0.075% yeast concentration was 532.33465 mg / 100g, for Fermented Glutinous Black Rice extract with 0.1% yeast concentration is 370.9936 mg / 100g, for Fermented Glutinous Black Rice extract with 0.125% yeast concentration is 198.995 mg / 100g, for Fermented Glutinous Black Rice extract with 0.15% yeast concentration is 189.11149 mg / 100g . The analysis showed that the higher the yeast concentration in fermented Fermented Glutinous Black Rice, the less anthocyanin levels contained in the sample. The effect of decreased levels of anthocyanin occurs due to the absorption of anthocyanin by the yeast cell wall. This is in accordance with Morata's research, 2016 concerning the Effect of Yeast on Khamr Color states that the yeast cell wall can adsorb anthocyanin pigments. Although the adsorption is only in a small color, it is reported that *S. Serevicae* chain can absorb anthocyanin in the range of 1.6 - 5.8% of anthocyanin levels. Besides that, the level of anthocyanin contained in Fermented Glutinous Black Rice which has undergone a fermentation process can decrease due to the heating stage in making Fermented Glutinous Black Rice. One factor that occurs during cooking is the development of starch grains from glutinous rice and rice which have an impact on the physical damage to the pericarp layer. Cooking causes the development of starch grains which have an impact on physical damage to the pericarp layer. The composition of starch which almost all consists of amylopectin causes Glutinous Black Rice does not expand in cooking (Mambrasar et al., 2010) In addition, statistical testing was carried out to determine the effect of adding yeast to anthocyanin levels. Obtained normality values so that the test is done with the Kruskal Wallis and Mann Whitney test. From the test results obtained that at the concentration of yeast 0.05%, 0.075%, 0.1% and 0.125% p value > 0.5 so that there are no significant differences.

Meanwhile, at a concentration of 0.15% the value of  $p < 0.5$ , which means that there is a meaningful difference,

or the concentration has a difference with other concentrations. DPPH trapping is often carried out for testing antioxidants from cysteine, glutathione, ascorbic acid, tocopherols, and the aromatic components of polyhydroxyl, olive oil, fruits, fruit juices and wine fermentation. The advantage of this method is that DPPH will act with the sample as a whole within a certain time and DPPH can react slowly even with weak antioxidants (Prakash, 2001). Antioxidant activity is related to rice type. Rice contains compounds such as tocopherol, oryzanol and polyphenols. Polyphenols such as vanillin, quercetin, and other phenolics are known to be potential antioxidants. Low molecular weight polysaccharides, proteins or peptides also affect antioxidants (Palaniveloo and Vairappan, 2013). Fermented Glutinous Black Rice and black rice are known to contain anthocyanin compounds which give a black color to the pericarp layer. In addition, the results obtained increase in the value of IC<sub>50</sub> with an increase in the concentration of yeast in the sample. Increased IC<sub>50</sub> illustrates a decrease in antioxidant activity. Because, the smaller the IC<sub>50</sub> value, the higher the free radical scavenging activity (Molyneux, 2004). This is like what happened in the study (Prabhu, 2014) the total antioxidant activity decreases in the presence of fermentation, because without the addition of oxygen, plant sterols quickly disrupt the yeast membrane's properties by becoming dominant dominant, causing slow fermentation. So during the 48th hour of fermentation many sterols are used by yeast so there is a decrease in total antioxidant activity.

## 5. Conclusion

Terdapat pengaruh konsentrasi ragi berupa penurunan hasil kadar antosianin dan aktivitas antioksidan pada setiap kenaikan konsentrasi ragi pada sampel. Didapatkan nilai kadar antosianin tertinggi sebesar 582.2344 mg/100g pada konsentrasi ragi 0,05% dan nilai aktivitas antioksidan tertinggi yang ditunjukkan dengan nilai IC<sub>50</sub> terendah sebesar 81,983 ppm pada konsentrasi ragi 0,05%.

## 6. Suggestion

Further research needs to be done to identify the type and quality of anthocyanin contained in Fermented Glutinous Black Rice more precisely such as using HPLC, so that the type and quality of anthocyanin can be known appropriately and it is necessary to improve the stability of the anthocyanin pigment contained in the Fermented Glutinous Black Rice so that it can be utilized as a source of natural dyes and herbal treatments.

## 7. Thank-you note

The researcher would like to thank to the Department of Pharmacy and the Department of Nutrition for the Poltekkes of the Ministry of Health in Bandung, as well as all parties involved in this research.

## 8. Reference

- [1] Agustinus, E.P., & Amran, H. 2009. Pembuatan Bioethanol dari Nira Siwalan Secara Fermentasi Fese Cair Menggunakan Fermipan. Semarang: Fakultas Teknik Jurusan Teknik Kimia Universitas Diponegoro
- [2] Aligita, Widhya. 2007. Skripsi. Isolasi Antosianin dari Ketan Hitam. Program Studi Sain dan Farmasi. Sekolah Farmasi Institut Teknologi Bandung.
- [3] Amarowicz, R., Naczki, M. & Shahidi, F. 2000. Antioxidant Activity of Crude Tannis of Canola And Rapeseed Hulls. *JAACS*, 77:95.
- [4] Basuki, N., Harijono, Kuswanto, & Damanhuri. 2005. Studi Pewarisan Antosianin pada Ubi Jalar Agravita, *Molecules*. 63-68
- [5] Benabadji SH, Wen R, Zheng JB, Dong XC, Yuan SG. 2004. Anticarcinogenic and Antioxidant Activity

of diindolylmethane Derivatives. *Journal Acta Pharmacologica Sinica*, 25 (5): 666-671.

[6] Bridle, P. dan Timberlake, C.F. 1997. Anthocyanin as Natural Food colours – Selected Aspects. *Food Chemistry*. Vol. 58 (1 – 2), pp 103 – 109

[7] Dykes, Rooney. 2007. Phenolic compounds in cereal grains and their health benefits, *Cereal Food Worlds*. Vol. 52.

[8] Elisa, P, Fulvio, M., & Johnson, C.S. 2013. The Case for Anthocyanin Consumption to Promote Human Health [A Review. *Comprehensive Reviews in Food Science and Food Safety*]. Vol 12.

[9] Fauziyah, N. 2015. Hubungan Konsumsi Tape Ketan Hitam Dengan Pencegahan Kejadian Sindroma Metabolik Pada Usia 40 Tahun Ke Atas Di Kabupaten Bandung Barat Provinsi Jawa Barat. Jakarta: Universitas Indonesia; Francis, F.J. 1982. Analysis of Anthocyanins. In P. Markakis (ed). *Anthocyanins as Food Colors*. New York: Academic press.

[10] Fauziyah, Roro Nur, Mimin Aminah, Osman Syarief, Holil M Par'I, Widi Hastuti, Surmita. 2020. Effectiveness of Steamed Brownies Base on Fermented Black Glutinous Rice on Decreased Waist Circumference in Abdominal Obesity. *Jurnal Ilmu dan Teknologi Kesehatan*. Vol x No x. Artikel <http://ejurnal.poltekkesjakarta3.ac.id/index.php/jitek>

[11] Fauziyah, Roro Nur, Putri, Mardiyah Maulida, Surmita. 2020. Effect of Pie Based on Fermented Black Glutinous Rice and Sweet Purple Potato to Frequency of Defecation in Adolescents with Constipation. *International Medical Journal*. Vol 25, Issue 04. Artikel <https://www.seroniijhou.com/volume/IMJ/25/4/effect-of-pie-based-on-fermented-black-glutinous-rice-and-sweet-purple-potato-to-frequency-of-defecation-in-adolescents-with-constipation-5ec80c6fdc2fa.pdf>

[12] Gandjar, Ibnu Gholib. 2007. *Kimia Farmasi Analisis*. Yogyakarta: Pustaka Pelajar

[13] Gary, F. 2007. God-given Foods Eating plan: for lifelong Health, Optimization of Hormones, Improved Athletic Performance. (<https://www.amazon.com/God-given-Foods-Eating-Plan-optimization/dp/1430319836>) diakses pada 11 Januari 2019

[14] Hadi, Spopyan., Thamrin., S, Moersidik, S.S., & Bahry, Syaiful. 2013. Karakteristik Dan Potensi Bioetanol Dari Nira Nipah Untuk Penerapan Skala Teknologi Tepat Guna. Pekanbaru: Program Studi Ilmu Lingkungan PPS Universitas Riau.

[15] Harborne. 2005. *Encyclopedia of Food and Color Additives*. New York: CRC Press Inc.,

[16] Hariadi, Arsyad. 2013. Prinsip Spektrofotometer-UV-VIS. Tanpa Kota Hidayat, Nur., Padaga, Masdiana., & Suhartini, Sri., 2006. *Mikrobiologi Industri*. Yogyakarta: C.V Andi Offset.

[17] Hutapea, E. R. F., Siahaan, L. O., & Tambun, R., 2014. Ekstraksi Pigmen Antosianin dari Kulit Rambutan (*Nephelium lappaceum*) dengan Pelarut Metanol. *Jurnal Teknik Kimia USU*, 3(2): 34-40

[18] Itani, T. & Ogawa, M. 2004. History and Recent Trends of Red Rice in Japan, *Nippon Sakumotsu Gakkai Kiji*. 73: 137-147.

- [19] Jackman R. L. dan J. L. Smith. 1996. Anthocyanins and Betalains. In Hendry.
- [20] G. A. P dan J. D. Houghton (eds). Natural Food Colorants, Second Edition. London: Chapman and Hall.
- [21] Jalil A.M, & Ismail A. 2008. Polyphenols in cocoa and cocoa product is there a link between antioxidant properties and health. *Molecules*. 13 (9): 2190-2219
- [22] Juliana, Reni., Julianti, Elisa., & Limbong, Lasma Nora. 2017. Pengaruh Metode dan Lama Fermentasi Terhadap Karakteristik Kimia Ubi Jalar Ungu. Medan
- [23] Karnajanawipagul, P, W. Nittayanuntawech, P. Rojsanga & L. Suntornsuk. 2010. Analysis of  $\beta$ - Carotene in Carrot by Spectrophotometry, *Mahidol University Journal of Pharmaceutical Science*. 37 (1-2), 8-16.
- [24] Khosman, A. 2002. Pangan dan Gizi untuk Kesehatan. Jakarta. PT Raja Grafindo Persada.
- [25] Kusmiyati, Mimin, Trinovani, Elvi, Fauziyah, Roro Nur. 2020. Activity of Mixed Ethanol Extract Selected Black Tea (*Camelia sinensis* L.) and Stevia (*Stevia rebaudiana* B.) as an Alternative Anti Diabetes Herbal Medicine. *International Medical Journal*. Vol 25, Issue 06. Artikel <https://www.seonijihou.com/volume/IMJ/25/6/activity-of-mixed-ethanol-extract-selected-black-tea-camelia-sinensis-l-and-stevia-stevia-rebaudiana-b-as-an-alternative-anti-diabetes-herbal-medicine-5ef522fdaa75e.pdf>
- [26] Lie, J., Zang, L., Yan, L., & Niu, L. 2012. Phenolic Compound and Antioxidan Activity of Bulb Extract of Six Liliun Species Native to China. *Molecules*. 17: 9361-9378.
- [27] Luciola S. 2012. Anthocyanins: Mechanism of action and therapeutic efficacy Research, *Signpost*.
- [28] Marzuki, Asnah. 2012. Kimia Analisis Farmasi. Makassar: Dua Satu Press
- [29] Mambrasar, Rinto Herry, Budhi Prasetyo, dan Martanto Martosupono, 2010. Antioksidan dan Imunomodulator Pada Serealia. UNS: Prosiding Seminar Nasional Pendidikan Biologi FKIP, 154-163.
- [30] Medina, Karina dkk. 2018. Effect of Non-Saccharomyces Yeast on Colour Anthocyanin, and Anthocyanin-Derived Pigments of Tannat Grapes during Fermentation. *American Society for Enology and Vitiiculture*. Amerika
- [31] Molyneux, P. 2004. The Use of Stable Free Radical Diphenylpicrylhydrazyl (DPPH) for Estimating Antioxidant Activity, *Journal Science Technologi*.
- [32] Morata, Antonio. 2016. Influence of Yeast in Wine Colour. INTECH. Madrid. Muchtadi D. 2013. Antioksidan dan Kiat Sehat di Usia Produktif. Bandung:Alfabeta;
- [33] Mudjajanto, E.S & L, N Yulianti. 2004. Membuat Aneka Roti. Jakarta: Penebar Swadaya.
- [34] Nur, M.A., M. Sjachri, dan K. Iskandarsyah. 1981. Kimia Dasar II. Bogor: Institut Pertanian Bogor Press. Palaniveloo, K dan Vairappan, C.S. 2013. Biochemical properties of rice wine produced from three different starter cultures. *Journal of Tropical Biology and Conservation* 10:31-41.

[35] Prabhu, Ashish. 2014. Effect of Yeast Fermentation on Nutraceutical and Antioxidant Properties of Rice Bran. *International journal of Agricultural and Food Science*. India. 4(1): 59-65

[36] Prakash A., 2001. Antioxidant Activity, Medaltion Laboratories Analitical Progres, Vol. 19 (2).

[37] Putranti. 2013. Skrining Fitokimia dan Aktivitas Antioksidan Ekstrak Rumput Laut *Sargassum duplicatum* dan *Turbinaria ornata* dari Jepara. Semarang: Fakultas Perikanan dan Ilmu Kelautan, Universitas Diponegoro.

[38] Rooney L.W., & Serna S. 2000. Shorgum. In Kulp, Karel., & Ponte, joseph. G.(Ed). *Handbook of Cereal Science and Technology*. New York : Marcel Dekker (<https://books.google.co.id/books?id=EUBZDwAAQBAJ&dq=Handbook+of+Cereal+Science+and+Technology.+rooney>) diakses pada 28 Januari 2019.

[39] Sandilova, E. Stintzing, F.C. & Carle, R. 2006. Thermal Degradation of Acylated and nonacylated Anthocyanins. *Journal of Food Science.I Jerman*.

[40] Sayuti, M., & Yenrina, R. 2015. *Antioksidan Alami dan Sintetik*. Padang: AU Press.

[41] Schneiter, R. 2004. *Genetics, Molecular and Cell Biology*. Swiss: Unversite De Fribourg Suisse.

[42] Seftian, Dedy., Antonius, Ferdinand., & Faizal, M. 2012. Pembuatan Etanol dari Kulit Pisang Menggunakan Metode Hidrolisis Enzimatik dan fermentasi. Palembang: Teknik Kimia Universitas Sriwijaya.

[43] Suzery, M., Lestari, S., & Cahyono, B. 2010. Penentuan Total Antosianin Dari Kelopak Bunga Rosela (*Hibiscus sabdariffa L*) dengan Metode Maserasi dan Sokshletasi, *Jurnal Sains dan Matematika*. 18(1): 1-6.

[44] Suhartatik, Nanik., & Karyantina, Merkuria. 2014. Karakteristik Fermentatis Medium de Mann Regosa Sharpe (MRS) Antosianin Beras ketan hitam (*oryza sativa var. glutinosa*) Menggunakan *Pediococcus pentosaceus* N.11.16. *Jurnal AgritechnSology*.

[45] Tamat, S. R., Wikanta, T., dan Maulina, L. S., 2007, Aktivitas Antioksidan dan Toksisitas Senyawa Bioaktif dari Ekstrak Rumput Laut Hijau *Ulva reticulate forsskal*, *Jurnal Ilmu Kefarmasian Indonesia*, 5(1): 31-36.

[46] Trinovani, Elvi, Afifah Riska Rafa, Fauziyah, Roro Nur. 2020. Determination of Antosianin Total Levels and Antioksidant Activities in Black Glutinous Rice Extract and Fermented Black Glutinous Extract. *International Medical Journal*. Vol 25, Issue 05. Artikel <https://www.seonijihou.com/volume/IMJ/25/5/determination-of-antosianin-total-levels-and-antioxidant-activities-in-black-glutinous-rice-extract-and-fermented-black-glutinous-rice-extract-5ec8099a8eb18.pdf>

[47] Vaughan D.A., Morishima H., & Kodawaki K. 2003. Diversity in the *Oryza* genus. *Current Opinion in Plant Biology*.

[48] Winarsi H. 2011. *Antioksidan Alami dan Radikal Bebas*. Yogyakarta: Kanisius. Yanuar W. 2009. *Aktivitas Antioksidan dan Imunomodulator Serealia*. Bogor. Institut Pertanian Bogor.



[49] Yustina, I. 2011. Studi Pengaruh Lama Fermentasi Tape Ketan Hitam terhadap Kadar Antosianin dan Aktivitas Antioksidan. Malang: Universitas Brawijaya.



This work is licensed under a Creative Commons Attribution Non-Commercial 4.0 International License.