

DAFTAR PUSTAKA

- Achadi, A. (2008) 'Regulasi Pengendalian Masalah Rokok di Indonesia', *Kesmas: National Public Health Journal*, 2(4), p. 161. doi: 10.21109/kesmas.v2i4.259.
- Afiani, Dina. (2019). Profil Nikotin, Kotinin dan Trans-3'-Hidroksikotinin di Urin Perokok Aktif. Repositori Institusi USU Departemen Pulmonologi dan Kedokteran Respirasi. Medan
- Aji, A., Maulinda, L. and Amin, S. (2015) 'Isolasi Nikotin Dari Puntung Rokok Sebagai Insektisida', *Jurnal Teknologi Kimia Unimal*, 4(1), pp. 100–120. <https://core.ac.uk/download/pdf/230118041.pdf>.
- Balhara YPS, Jain R. (2013). A receiver operated curve-based evaluation of change in sensitivity and specificity of cotinine urinalysis for detecting active tobacco use. *Journal of Cancer Research and Therapeutics*. 9(1):84-89.
- Benowitz, N.L. (2009). Pharmacology of nicotine: Addiction, smoking-induced disease, and therapeutics, *National Institute of Health*, vol.49, pp: 57-71. <https://www.annualreviews.org/doi/full/10.1146/annurev.pharmtox.48.113006.094742#article-denial> diakses pada tanggal 19 September 2020
- Benowitz, N.L., (2010). Nicotine Addictiome. *N Eng. J Med*. 362 (24): 2295–303.
- Bustan, M.N. (2007). Epidemiologi Penyakit Tidak Menular. Jakarta: PT Rineka Cipta
- Elmanfe, G. and Omar, S. (2020) 'Determination of Nicotine in Libyan Smokers and Nonsmokers' Urine samples by Ultraviolet-Visible Spectrophotometry, *Journal of Pure & Applied Sciences*. Chemistry Department, Faculty of Science, Omar Al-Mukhtar University, El-Bieda, Libya
- Hans Tendra. (2003). Merokok dan Kesehatan. Jakarta: Rineka Cipta
- Hukkanen, J., Jacob, P., Benowitz, N.L. (2005). Metabolism and disposition kinetics of nicotine, *American Society for Pharmacology and Experimental Therapeutics*, vol. 57, pp: 79-115.
- Husaini, A. (2007). Tobat Merokok. Jakarta: Pustaka Iman.

- Inayatillah, I. R., Syahrudin, E. and Susanto, A. D. (2014) 'Kadar Karbon Monoksida Udara Ekspirasi pada Perokok dan Bukan Perokok serta Faktor-Faktor yang Mempengaruhi Exhaled Carbon Monoxide Levels in Smokers and Non Smokers and Related Factors', *Jurnal Respirologi Indonesia*, 34(4), pp. 180–90. Available at: <http://jurnalrespirologi.org/wp-content/uploads/2015/08/JRI-Oct-2014-34-4-180-90.pdf>.
- Ispranoto, Tri. (2017) Ada Larangan, Perokok di Cijawura Bandung Tak Sampai 100 Orang. <https://news.detik.com/berita-jawa-barat/d-3658955/ada-larangan-perokok-di-cijawura-bandung-tak-sampai-100-orang> diakses pada tanggal 22 Maret 2021
- Jacob, P., Yu, L., Duan, M., Ramos, L., Yturralde, O., Benowitz, N.L. (2011). Determination of the nicotine metabolites cotinine and trans-3-hydroxycotinine in biologic fluids of smokers and non-smokers using liquid chromatography-tandem mass spectrometry: Biomarkers for tobacco smoke exposure and for phenotyping cytochrome P450 2A6 activity. *J Chromatogr B*. 879:267-76.
- Jaya, M., (2009). Pembunuh Berbahaya Itu Bernama Rokok. 1st ed. Yogyakarta: Riz'ma
- Kawasaki, Y. *et al.* (2020) 'Effects of smoking cessation on biological monitoring markers in urine', *Genes and Environment*. 42(1), pp. 1–9. doi: 10.1186/s41021-020-00165-z.
- Kulza M, Wozniak A, Przybylowska SM, Czarnywojtek A, Kurhanska-Flisykowska A, Florek E. (2012). Saliva cotinine determination using high-performance liquid chromatography with diode-array detection. *Przegl Lek*. 69(10):837-40.
- Lianzi, I. and Pitaloka, E. (2014) 'Hubungan pengetahuan tentang rokok dan perilaku merokok pada staf administrasi universitas esa unggul', *Jurnal Inohim*, 2(1), pp. 67–81.
- Masturoh, I., dan N. Anggita. (2018). Metodologi Penelitian Kesehatan. Kementerian Kesehatan RI. Jakarta.
- Matsumoto, A., Ichiba, M., Payton, N. M., Oishi, H., & Hara, M. (2013). Simultaneous measurement of urinary total nicotine and cotinine as biomarkers of active and passive smoking among Japanese individuals. *Environmental health and preventive medicine*, 18(3), 244–250. <https://doi.org/10.1007/s12199-012-0307-5>
- Mitali, R., *et al.* (2016). Diagnostic Methods for Detection of Cotinine Level in Tobacco Users: A Review. *Clinic and Diagnosis Respiratory Journal*. 10(3)

- Monique, As. (2005). *Menghindari Merokok*. Jakarta: PT Musi Perkasa Utama
- National Center for Biotechnology Information (2021). PubChem Compound Summary for CID 89594, Nicotine. <https://pubchem.ncbi.nlm.nih.gov/compound/Nicotine> diakses pada tanggal 12 Maret 2021
- Neal, L.B., Janne, H., and Peyton, J. (2009). Nicotine Chemistry, Metabolism, Kinetics and Biomarkers. *Handb Exp Pharmacol.* (192): 29–60
- Nima, B., *et al.* (2014). Correlation between Cigarette Smoking and Urine Cotinine Level in Gastric Cancer Patients. *Iranian Journal of Pharmaceutical Research.* 13 (1): 313-318
- Parker DR, Thomas M, Richard W, Jeff W, Upegui DI, Heimdal J. (2002). The accuracy of self-reported smoking status assessed by cotinine test strips. *Nicotine & Tobacco Research.* 4:305-09.
- Patramurti, C. *et al.* (2017). Inter-Individual Variability of Cytochrome P450 2A6 Activity in Javanese Smokers Urine. *Media Penelitian dan Pengembangan Kesehatan,* 27(3), pp. 133–140. doi: 10.22435/mpk.v27i3.4777.133-140.
- Peraturan Pemerintah Republik Indonesia nomor 81 tahun 1999 tentang Pengamanan Rokok Bagi Kesehatan
- Peraturan Walkota Bandung No.315 tahun 2017 tentang Kawasan Tanpa Rokok (KTR)
- Pitriyanti, Luh. (2018). Proporsi dan Faktor-Faktor yang Mempengaruhi Penggunaan Rokok Elektronik Pada Mahasiswa Universitas Udayana: Dengan Landasan Social Cognitive Theory. Fakultas Kedokteran Universitas Udayana Denpasar.
- Pusat Data dan Informasi Kementerian Kesehatan RI. (2019). Situasi Umum Konsumsi Tembakau di Indonesia (Current Tobacco Consumption in Indonesia).
- Raja, M. (2016) ‘Diagnostic Methods for Detection of Cotinine Level in Tobacco Users: A Review’, *Journal of Clinical and Diagnostic Research,* (March). doi: 10.7860/jcdr/2016/17360.7423.
- Ramdhani, Dendi. (2017). Warga Pandanwangi Bandung Sukses Ciptakan Kawasan Tanpa Rokok.
<https://regional.kompas.com/read/2017/09/27/06540011/warga-pandanwangi-bandung-sukses-ciptakan-kawasan-tanpa-rokok?page=all>. diakses pada tanggal 22 Maret 2021

- Rigotti, N. A., Pipe, A. L., Benowitz, N. L., Arteaga, C., Garza, D., & Tonstad, S. (2010). Efficacy and safety of varenicline for smoking cessation in patients with cardiovascular disease: a randomized trial. *Circulation*, *121*(2), 221–229. <https://doi.org/10.1161/CIRCULATIONAHA.109.869008>
- Riset Kesehatan Dasar (Riskesdas). (2019). Badan Penelitian dan Pengembangan Kesehatan Kementerian RI tahun 2018
- Sanner, Tore. Grimsrud, Tom K. (2015). Nicotine: Carcinogenicity And Effects On Response To Cancer Treatment – A Review. USA : Volume 5 | Article 196.
- Sitepoe. (2000). Kekhususan Rokok Indonesia. Jakarta: PT Gramedia Widiasarana Indonesia.
- Susanna, D., Hartono, B., Fauzan, H. (2003). Penentuan kadar nikotin dalam asap rokok. *Jurnal Universitas Indonesia*. Jakarta: *Makara Kesehatan*;7.
- Tang EA, Delaimy WKA, Ashley DL, Benowitz N, Bernert JT, Kim S et al. (2003). Assessing second hand smoke using biological markers. *Tobacco control*. *22*(3):164-71.
- Urin, P. *et al.* (2014) ‘Gangguan Fungsi Paru Dan Kadar Cotinine Pada Urin Karyawan Yang Terpapar Asap Rokok Orang Lain’, *KEMAS: Jurnal Kesehatan Masyarakat*, *10*(1), pp. 43–52. doi: 10.15294/kemas.v10i1.3069.
- Vusvitasari, R., Nugroho, S. and Akbar, S. (2016) ‘Kajian Hubungan Koefisien Korelasi Pearson (ρ), Spearman-’, *Journal Statistika*, pp. 41–54.
- World Health Organization. (2015). Global Youth Tobacco Survey Indonesia Report, 2014. Jakarta: WHO
- World Health Organization. (2018). WHO Report on The Global Tobacco. https://www.who.int/health-topics/tobacco#tab=tab_1 diakses pada tanggal 19 September 2020
- World Health Organization. (2021). The importance of tobacco cessation in the context of the COVID-19 pandemic <https://www.who.int/news-room/events/detail/2021/03/16/default-calendar/the-importance-of-tobacco-cessation-in-the-context-of-the-COVID-19-pandemic> diakses pada tanggal 17 Maret 2021
- Yildiz, D. (2004). Nicotine, its metabolism and an overview of its biological effects, *Elsevier*, pp: 619-632

Zhu, A.Z.X., Binnington, A.J., Renner, C.C., Lanier, A.P., Hatsukami, D.K., Stepanov, I., *et al.* (2013). Alaska native smokers and smokeless tobacco users with slower CYP2A6 activity have lower tobacco consumption, lower tobacco specific nitrosamine exposure and lower tobacco-specific nitrosamine bioactivation, *Carcinogenesis*, vol.34, no.1, pp: 93-101.